

STEIN'S METHOD FOR STEADY-STATE DIFFUSION APPROXIMATIONS: AN INTRODUCTION THROUGH THE ERLANG-A AND ERLANG-C MODELS

BY ANTON BRAVERMAN J.G. DAI AND JIEKUN FENG

Cornell University

This paper provides an introduction to the Stein method framework in the context of steady-state diffusion approximations. The framework consists of three components: the Poisson equation and gradient bounds, generator coupling, and moment bounds. Working in the setting of the Erlang-A and Erlang-C models, we prove that both Wasserstein and Kolmogorov distances between the stationary distribution of a normalized customer count process, and that of an appropriately defined diffusion process decrease at a rate of $1/\sqrt{R}$, where R is the offered load. Furthermore, these error bounds are *universal*, valid in any load condition from lightly loaded to heavily loaded. In the context of the Erlang-C model, we find that the diffusion approximation of higher moments is *not universal*.

1. Introduction. In [4], the authors developed a framework based on Stein's method [15, 16] to prove the rates of convergence for steady-state diffusion approximations. Using their framework, they were able to prove convergence rates for the steady-state approximation of the $M/Ph/n + M$ system, the many server queue with customer abandonment and phase-type service times, in the Halfin-Whitt regime [11]. The framework in [4] is modular and has four components: the Poisson equation and gradient bounds, generator coupling, moment bounds, and state space collapse (SSC). The purpose of this paper is to provide an accessible introduction to the Stein framework, focusing on two simple and yet fundamental systems. They are the $M/M/n + M$ system, known as the Erlang-A system, and $M/M/n$ system, known as the Erlang-C system. The accessibility is due to the fact that both systems can be represented by a one-dimensional continuous time Markov chain (CTMC). In addition, by focusing on these two systems we are able to present some sharp results that serve as benchmarks that future research should aspire to meet.

Stein's method is a powerful method that has been widely used in probability, statistics, and their wide range of applications such as bioinformatics; see, for example, the survey papers [5, 14], the recent book [6] and the references within. Barbour [2] has used Stein's method to establish convergence rates for approximations by a diffusion process.

Both Erlang-A and Erlang-C systems have n homogeneous servers that serve customers

MSC 2010 subject classifications: Primary 60K25; secondary 60F99, 60J60

Keywords and phrases: Stein's method, steady-state, diffusion approximation, convergence rates, Erlang-A, Erlang-C

in a first-come-first-serve manner. Customers arrive according to a Poisson process with rate λ , and customer service times are assumed to be i.i.d. having exponential distribution with mean $1/\mu$. In the Erlang-A system, each customer has a patience time and when his waiting time in queue exceeds his patience time, he abandons the queue without service; the patience times are assumed to be i.i.d. having exponential distribution with mean $1/\alpha$. The Erlang-A system is a special case of the systems studied in [4], where SSC played an important role. The systems in this paper can be represented by a one-dimensional CTMC, meaning that there is no need to invoke SSC. Therefore, this paper illustrates only the first three components of the framework proposed in [4].

We will study the birth-death process

$$(1.1) \quad X = \{X(t), t \geq 0\},$$

where $X(t)$ is the number of customers in the system at time t . In the Erlang-A system, α is assumed to be positive and therefore the mean patience time is finite. This guarantees that the CTMC X is positive recurrent. In the Erlang-C system, $\alpha = 0$, and in order for the CTMC to be positive recurrent we need to assume that the offered load to the system, defined as $R = \lambda/\mu$, satisfies

$$(1.2) \quad R < n.$$

For both Erlang-A and Erlang-C systems, we use $X(\infty)$ to denote the random variable having the stationary distribution of X .

The following theorem illustrates the type of result that can be obtained by Stein's method. Consider the case when $\alpha = 0$ and (1.2) is satisfied. Set $\tilde{X}(\infty) = (X(\infty) - R)/\sqrt{R}$, and let $Y(\infty)$ denote a continuous random variable on $\mathbb{R}$ having density

$$(1.3) \quad \kappa \exp\left(\frac{1}{\mu} \int_0^x b(y) dy\right),$$

where $\kappa > 0$ is a normalizing constant that makes the density integrate to one,

$$(1.4) \quad b(x) = [(x + \zeta)^- - \zeta^-] \mu, \quad \text{and} \quad \zeta = (R - n)/\sqrt{R}.$$

The following theorem is an example of our main results, the rest of which are stated in Section 2. The framework to prove this theorem is developed in Section 3, where we describe each component of the Stein framework.

THEOREM 1. *Consider the Erlang-C system ($\alpha = 0$). For all $n \geq 1$, $\lambda > 0$, and $\mu > 0$ satisfying $1 \leq R < n$,*

$$(1.5) \quad d_W(\tilde{X}(\infty), Y(\infty)) \equiv \sup_{h \in \text{Lip}(1)} |\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))| \leq \frac{205}{\sqrt{R}},$$

where

$$\text{Lip}(1) = \{h : \mathbb{R} \rightarrow \mathbb{R}, |h(x) - h(y)| \leq |x - y|\}.$$

$n = 5$			$n = 500$		
R	$\mathbb{E}X(\infty)$	Error	R	$\mathbb{E}X(\infty)$	Error
3	3.35	0.10	300	300.00	6×10^{-14}
4	6.22	0.20	400	400.00	2×10^{-6}
4.9	51.47	0.28	490	516.79	0.24
4.95	101.48	0.29	495	569.15	0.28
4.99	501.49	0.29	499	970.89	0.32

TABLE 1

Comparing the error $|\mathbb{E}X(\infty) - (R + \sqrt{R}\mathbb{E}Y(\infty))|$ for different system configurations.

The framework developed in [4] was inspired largely by the work of Gurvich in [9] who developed methodologies to prove statements similar to Theorem 1 for a broad class of multidimensional CTMCs. Along the way, he independently rediscovered many of the ideas central to Stein's method in the setting of steady-state diffusion approximations. See [4] for a more detailed discussion of Gurvich's results.

Several points are worth mentioning. First, we note that Theorem 1 is not a limit theorem. Steady-state approximations are usually justified by some kind of limit theorem. That is, one considers a sequence of queueing systems and proves that the corresponding sequence of steady-state distributions converges to some limiting distribution as traffic intensity approaches one, or as the number of servers goes to infinity. In contrast, our theorem holds for any finite parameter choices of λ, n , and μ satisfying (1.2) and $R \geq 1$. Second, the error bound in (1.5) is *universal*, as it does not assume any relationship between λ, n , and μ , other than the stability condition (1.2) and the condition that $R \geq 1$. The latter condition is a mere convenience, as Theorem 1 could be restated without it, but the error bound would then contain some terms involving $1/R$. In particular, universality means that the error bound holds when λ, n , and μ satisfy one of the following conditions:

$$\begin{aligned} n &= \lceil R + \beta R \rceil, \quad \beta > 0, \\ n &= \lceil R + \beta \sqrt{R} \rceil, \quad \beta > 0, \\ n &= \lceil R + \beta \rceil, \quad \beta > 0. \end{aligned}$$

The first two parameter regimes above describe the quality-driven (QD), and quality-and-efficiency-driven (QED) regimes, respectively. The last regime is the nondegenerate-slowdown (NDS) regime, which was studied in [1, 18]. Universal approximations were previously studied in [10, 17]. Third, as part of the universality of Theorem 1, we see that

$$(1.6) \quad |\mathbb{E}X(\infty) - (R + \sqrt{R}\mathbb{E}Y(\infty))| \leq 205.$$

For a fixed n , let $\rho = R/n \uparrow 1$. One expects that $\mathbb{E}X(\infty)$ be on the order of $1/(1 - \rho)$. Conventional heavy-traffic limit theorems often guarantee that the left hand side of (1.6) is at most $o(1/\sqrt{1 - \rho})$, whereas our error is bounded by a constant regardless of the load condition. This suggests that the diffusion approximation for the Erlang-C system is

R	$\mathbb{E}(\tilde{X}(\infty))^2$	$ \mathbb{E}(\tilde{X}(\infty))^2 - \mathbb{E}(Y(\infty))^2 $	$\mathbb{E}(\tilde{X}(\infty))^{10}$	$ \mathbb{E}(\tilde{X}(\infty))^{10} - \mathbb{E}(Y(\infty))^{10} $
300	1	4.55×10^{-15}	9.77×10^2	31.58
400	1	5.95×10^{-7}	9.70×10^2	24.44
490	6.96	0.11	7.51×10^9	7.01×10^8
495	31.56	0.27	9.10×10^{12}	4.34×10^{11}
499	9.47×10^2	1.59	1.07×10^{20}	1.03×10^{18}
499.9	9.94×10^4	16.50	1.13×10^{30}	1.09×10^{27}

TABLE 2

Approximating the second and tenth moments of $\tilde{X}(\infty)$ with $n = 500$. The approximation error grows as R approaches n and suggests that the diffusion approximation of higher moments is not universal.

accurate not only as $R \rightarrow \infty$, but also in the heavy-traffic setting when $R \rightarrow n$. Table 1 contains some numerical results where we calculate the error on the left side of (1.6). The constant 205 in (1.6) is unlikely to be a sharp upper bound. In this paper we did not focus too much on optimizing the upper bound, as Stein's method is not known for producing sharp constants.

From Theorem 1 we know that the first moment of $\tilde{X}(\infty)$ can be approximated universally by the first moment of $Y(\infty)$. It is natural to ask what can be said about the approximation of higher moments. We performed some numerical experiments in which we approximate the second and tenth moments of $\tilde{X}(\infty)$ in a system with $n = 500$. The results are displayed in Table 2. One can see that the approximation errors grow as the offered load R gets closer to n , suggesting that the approximation of higher moments of $\tilde{X}(\infty)$ is *not universal*. This differs from the approximation errors of the first moment in Table 1, where the errors approached some limiting value. We explore this topic further in Section 6, where Theorem 5 presents error bounds for approximating higher moments.

Theorem 1 provides rates of convergence under the Wasserstein metric [14]. The Wasserstein metric is one of the most commonly studied metrics in the context of Stein's method. This is because the space $\text{Lip}(1)$ is relatively simple to work with, but is also rich enough so that convergence under the Wasserstein metric implies the convergence in distribution [7]. Another metric commonly studied in problems involving Stein's method is the Kolmogorov metric, which measures the distance between cumulative distribution functions of two random variables. The Kolmogorov distance between $\tilde{X}(\infty)$ and $Y(\infty)$ is

$$\sup_{h \in \mathcal{H}_K} |\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))|, \quad \text{where} \quad \mathcal{H}_K = \{1_{(-\infty, a]}(x) : a \in \mathbb{R}\}.$$

Theorems 3 and 4 of Section 2 involve the Kolmogorov metric. A general trend in Stein's method is that establishing convergence rates for the Kolmogorov metric often requires much more effort than establishing rates for the Wasserstein metric, and our problem is no exception. The extra difficulty always comes from the fact that the test functions belonging to the class $\mathcal{H}_K$ are discontinuous, whereas the ones in $\text{Lip}(1)$ are Lipschitz-continuous. In Section 5, we describe how to overcome this difficulty in our model setting.

The first paper to have established convergence rates for steady-state diffusion approxi-

mations was [10], which studied the Erlang-A system using an excursion based approach. Their approximation error bounds are universal. Although the authors in [10] did not study the Erlang-C system, their approach appears to be extendable to it as well. However, their method is not readily generalizable to the multi-dimensional setting. In this paper we also derive approximation error bounds for the Erlang-A model by using Stein's method. Our notion of universality is stronger than the one in [10]. In most of their results, the authors of [10] allowed λ and n to vary without restriction, but they fix μ and α . The only exception is that they consider separately the NDS regime when $\mu = \mu(\lambda) = \beta\sqrt{\lambda}$ and $\lambda = n\mu + \beta_1\mu$ for some $\beta > 0$ and $\beta_1 \in \mathbb{R}$. In contrast, our error bound only depends on the ratio α/μ ; see Theorems 2 and 4.

We wish to point out that the results proved in this paper are quite sharp, and proving analogous results in a high dimensional setting is likely much more difficult. For instance, the constants in the error bounds in each theorem of this paper can be recovered explicitly, but this is not true in the problem studied in [4]. Moreover, the result of [4] is restricted to the Halfin–Whitt regime, and is not universal. All of this is because the model considered there is high dimensional.

1.1. *Notation.* For two random variables U and V , define their Wasserstein distance to be

$$(1.7) \quad d_W(U, V) = \sup_{h \in \text{Lip}(1)} |\mathbb{E}[h(U)] - \mathbb{E}[h(V)]|,$$

where

$$\text{Lip}(1) = \{h : \mathbb{R} \rightarrow \mathbb{R}, |h(x) - h(y)| \leq |x - y|\}.$$

It is known, see for example [14], convergence under the Wasserstein metric implies convergence in distribution. When $\text{Lip}(1)$ in (1.7) is replaced by

$$(1.8) \quad \mathcal{H}_K = \{1_{(-\infty, a]}(x) : a \in \mathbb{R}\},$$

the corresponding distance is the Kolmogorov distance, denoted by $d_K(U, V)$. When V has a density that is bounded by $C > 0$, then [14, Proposition 1.2] implies that

$$(1.9) \quad d_K(U, V) \leq \sqrt{2Cd_W(U, V)}.$$

For $a, b \in \mathbb{R}$, we use $a^+, a^-, a \wedge b$, and $a \vee b$ to denote $\max(a, 0)$, $\max(-a, 0)$, $\min(a, b)$, and $\max(a, b)$, respectively.

The rest of the paper is structured as follows. In Section 2 we state our main results. In Section 3 we present the Stein framework needed to prove our main results, Theorems 1–4, by introducing three ingredients central to our framework. Namely, the Poisson equation and gradient bounds, generator coupling, and moment bounds. In Section 4 we prove Theorems 1 and 2, which deal with the Wasserstein metric. The Kolmogorov metric presents an additional challenge, because the test functions are discontinuous. In Section 5 we address this new challenge and then prove Theorems 3 and 4. In Section 6, we discuss the non-universality of approximating higher moments in the Erlang-C model.

2. Main results. Recall the offered load $R = \lambda/\mu$. For notational convenience we define $\delta > 0$ as

$$(2.1) \quad \delta = \frac{1}{\sqrt{R}} = \sqrt{\frac{\mu}{\lambda}}.$$

Let $x(\infty)$ be the unique solution to the flow balance equation

$$(2.2) \quad \lambda = (x(\infty) \wedge n)\mu + (x(\infty) - n)^+ \alpha.$$

Here, $x(\infty)$ is interpreted as the equilibrium number of customers in the corresponding fluid model, and is the point at which the arrival rate equals the departure rate. The latter is the sum of the service completion rate and the customer abandonment rate with $x(\infty)$ customers in the system. One can check that the flow balance equation has a unique solution $x(\infty)$ given by

$$(2.3) \quad x(\infty) = \begin{cases} n + \frac{\lambda - n\mu}{\alpha} & \text{if } R \geq n, \\ R & \text{if } R < n. \end{cases}$$

By noting that the number of busy servers $x(\infty) \wedge n$ equals n minus the number of idle servers $(x(\infty) - n)^-$, the equation in (2.2) becomes

$$(2.4) \quad \lambda - n\mu = (x(\infty) - n)^+ \alpha - (x(\infty) - n)^- \mu.$$

We note that $x(\infty)$ is well-defined even when $\alpha = 0$, because in that case we always assume that $R < n$.

We consider the CTMC

$$(2.5) \quad \tilde{X} = \{\tilde{X}(t) \equiv \delta(X(t) - x(\infty)), t \geq 0\},$$

and let the random variable $\tilde{X}(\infty)$ have its stationary distribution. Define

$$(2.6) \quad \zeta = \delta(x(\infty) - n),$$

and

$$(2.7) \quad b(x) = [(x + \zeta)^- - \zeta^-] \mu - [(x + \zeta)^+ - \zeta^+] \alpha \quad \text{for } x \in \mathbb{R},$$

with convention that α is set to be zero in the Erlang-C system. Throughout this paper, let $Y(\infty)$ denote a continuous random variable on $\mathbb{R}$ having density

$$(2.8) \quad \nu(x) = \kappa \exp\left(\frac{1}{\mu} \int_0^x b(y) dy\right),$$

where $\kappa > 0$ is a normalizing constant that makes the density integrate to one. Note that these definitions are consistent with (1.3) and (1.4).

THEOREM 2. *Consider the Erlang-A system ($\alpha > 0$). There exists an increasing function $C_W : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that for all $n \geq 1, \lambda > 0, \mu > 0$, and $\alpha > 0$ satisfying $R \geq 1$,*

$$(2.9) \quad d_W(\tilde{X}(\infty), Y(\infty)) \leq C_W(\alpha/\mu)\delta.$$

REMARK 1. The proof of Theorems 1 and 2 uses the same ideas. Therefore, for the sake of brevity, we only give an outline for the proof of Theorem 2 and its ingredients without filling in all the details. It is for this reason that we do not write out the explicit form of $C_W(\alpha/\mu)$, although it can be obtained from the proof. The same is true for Theorem 4 below.

Although (1.9) often implies Kolmogorov metric convergence from the Wasserstein metric convergence, the constant C in (1.9) may depend on λ, n, μ , and α in general. Furthermore, the upper bound on the Kolmogorov metric implied by (1.9) is usually suboptimal, in the sense that the order of magnitude of the bound is incorrect. Therefore, for the bounds on the Kolmogorov metric we need to prove separately the following two theorems.

THEOREM 3. *Consider the Erlang-C system ($\alpha = 0$). For all $n \geq 1, \lambda > 0$, and $\mu > 0$ satisfying $1 \leq R < n$,*

$$(2.10) \quad d_K(\tilde{X}(\infty), Y(\infty)) \leq 188\delta.$$

THEOREM 4. *Consider the Erlang-A system ($\alpha > 0$). There exists an increasing function $C_K : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that for all $n \geq 1, \lambda > 0, \mu > 0$, and $\alpha > 0$ satisfying $R \geq 1$,*

$$(2.11) \quad d_K(\tilde{X}(\infty), Y(\infty)) \leq C_K(\alpha/\mu)\delta.$$

A version of Theorems 2 and 4 was first proved in the pioneering paper [10] using an excursion based approach. Theorems 1 and 3 are new.

3. Outline of the Stein Framework. In this section we introduce the main tools needed to prove Theorems 1–4. At this point we do not restrict ourselves to either the Erlang-A or Erlang-C systems, as the framework outlined here is generic and holds for both systems.

The following is an informal outline of the rest of this section. We know that $\tilde{X}(\infty)$ follows the stationary distribution of the CTMC $\tilde{X}$, and that this CTMC has a generator $G_{\tilde{X}}$. To $Y(\infty)$, we will associate a diffusion process with generator G_Y . We will start by fixing a test function $h : \mathbb{R} \rightarrow \mathbb{R}$ and deriving the identity

$$(3.1) \quad |\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))| = |\mathbb{E}G_{\tilde{X}}f_h(\tilde{X}(\infty)) - \mathbb{E}G_Yf_h(Y(\infty))|,$$

where $f_h(x)$ is a solution to the Poisson equation

$$G_Y f_h(x) = \mathbb{E}h(Y(\infty)) - h(x), \quad x \in \mathbb{R}.$$

We then focus on bounding the right hand side of (3.1), which is easier to handle than the left hand side. This is done by performing a Taylor expansion of $G_{\tilde{X}} f_h(x)$ in Section 3.3. To bound the error term from the Taylor expansion, we require bounds on various moments of $|\tilde{X}(\infty)|$, as well as the derivatives of $f_h(x)$. We refer to the former as moment bounds, and the latter as gradient bounds. These are presented in Sections 3.4 and 3.5, respectively.

3.1. The Poisson Equation of a Diffusion Process. The random variable $Y(\infty)$ in Theorems 1–4 is well-defined and its density is given in (2.8). It turns out that $Y(\infty)$ has the stationary distribution of a diffusion process $Y = \{Y(t), t \geq 0\}$, which we will define shortly. We do not prove this claim in this paper since it is not used anywhere in this paper. Nevertheless, it is helpful to think of $Y(\infty)$ in the context of diffusion processes. The diffusion process Y is the one-dimensional piecewise Ornstein–Uhlenbeck (OU) process. Its generator is given by

$$(3.2) \quad G_Y f(x) = b(x)f'(x) + \mu f''(x) \quad \text{for } x \in \mathbb{R}, f \in C^2(\mathbb{R}),$$

where $b(x)$ is defined in (2.7). Clearly, $b(0) = 0$, and $b(x)$ is Lipschitz continuous. Indeed,

$$|b(x) - b(y)| \leq (\alpha \vee \mu) |x - y| \quad \text{for } x, y \in \mathbb{R}.$$

Since the diffusion process Y depends on parameters λ, n, μ , and α in an arbitrary way, there is no appropriate way to talk about the limit of $Y(\infty)$ in terms of these parameters. Therefore, we call Y a *diffusion model*, as opposed to a *diffusion limit*. Having a diffusion model whose input parameters are directly taken from the corresponding Markov chain model is critical to achieve *universal accuracy*. In other words, this diffusion model is accurate in any parameter regime, from underloaded, to critically loaded, and to overloaded.

The main tool we use is known as the Poisson equation. It allows us to say that $Y(\infty)$ is a good estimate for $\tilde{X}(\infty)$ if the generator of Y behaves similarly to the generator of $\tilde{X}$, where $\tilde{X}$ is defined in (2.5). Let $\mathcal{H}$ be a class of functions $h : \mathbb{R} \rightarrow \mathbb{R}$, to be specified shortly. For each function $h(x) \in \mathcal{H}$, consider the Poisson equation

$$(3.3) \quad G_Y f_h(x) = \mu f_h''(x) + b(x)f_h'(x) = \mathbb{E}h(Y(\infty)) - h(x), \quad x \in \mathbb{R}.$$

One may verify by differentiation that for all functions $h : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $\mathbb{E}|h(Y(\infty))| < \infty$, the Poisson equation has a family of solutions of the form

$$(3.4) \quad f_h(x) = a_1 + \int_0^x \left[a_2 \frac{1}{\nu(u)} + \frac{1}{\nu(u)} \int_{-\infty}^u \frac{1}{\mu} (\mathbb{E}h(Y(\infty)) - h(y)) \nu(y) dy \right] du,$$

where $a_1, a_2 \in \mathbb{R}$ are arbitrary constants, and $\nu(x)$ is as in (2.8).

In this paper, we take $\mathcal{H} = \text{Lip}(1)$ when we deal with the Wasserstein metric (Theorems 1 and 2), and we choose $\mathcal{H} = \mathcal{H}_K$ when we deal with the Kolmogorov metric (Theorems 3 and 4). We claim that that $|\mathbb{E}h(Y(\infty))| < \infty$. Indeed, when $\mathcal{H} = \mathcal{H}_K$, this clearly holds. When

$\mathcal{H} = \text{Lip}(1)$, without loss of generality we take $h(0) = 0$ in (3.3), and use the Lipschitz property of $h(x)$ to see that

$$|\mathbb{E}h(Y(\infty))| \leq \mathbb{E}|Y(\infty)| < \infty,$$

where the finiteness of $\mathbb{E}|Y(\infty)|$ will be proved in (C.16).

From (3.3), one has

$$(3.5) \quad |\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))| = |\mathbb{E}G_Y f_h(\tilde{X}(\infty))|.$$

In (3.5), $\tilde{X}(\infty)$ has the stationary distribution of the CTMC $\tilde{X}$, not necessarily defined on the same probability space of $Y(\infty)$. Actually, $\tilde{X}(\infty)$ in (3.5) can be replaced by any other random variable, although one does not expect the error on the right side to be small if this random variable has no relationship with the diffusion process Y .

3.2. Comparing Generators. To prove Theorems 1–4, we need to bound the right side of (3.5). The CTMC $\tilde{X}$ defined in (2.5) also has a generator. We bound the right side of (3.5) by showing that the diffusion generator in (3.2) is similar to the CTMC generator.

For any $k \in \mathbb{Z}_+$, we define $x = x_k = \delta(k - x(\infty))$. Then for any function $f : \mathbb{R} \rightarrow \mathbb{R}$, the generator of $\tilde{X}$ is given by

$$(3.6) \quad G_{\tilde{X}} f(x) = \lambda(f(x + \delta) - f(x)) + d(k)(f(x - \delta) - f(x)),$$

where

$$(3.7) \quad d(k) = \mu(k \wedge n) + \alpha(k - n)^+,$$

is the departure rate corresponding to the system having k customers. One may check that

$$(3.8) \quad b(x) = \delta(\lambda - d(k)).$$

The relationship between $G_{\tilde{X}}$ and the stationary distribution of $\tilde{X}$ is illustrated by the following lemma.

LEMMA 1. *Let $f(x) : \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $|f(x)| \leq C(1+x)^2$ for some $C > 0$ (i.e. $f(x)$ is dominated by a quadratic function), and assume that the CTMC $\tilde{X}$ is positive recurrent. Then*

$$\mathbb{E}[G_{\tilde{X}} f(\tilde{X}(\infty))] = 0.$$

REMARK 2. As we will see in Lemmas 4–7, there is a family of solutions to the Poisson equation (3.3) whose first derivatives grow at most linearly in both the Wasserstein and Kolmogorov settings, meaning that these solutions satisfy the conditions of Lemma 1.

The proof of Lemma 1 is provided in Section A, and relies on Proposition 3 of [8]. Suppose for now that for any $h(x) \in \mathcal{H}$, the solution to the Poisson equation $f_h(x)$ satisfies the conditions of Lemma 1. We can apply Lemma 1 to (3.5) to see that

$$\begin{aligned}
 |\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))| &= |\mathbb{E}G_Y f_h(\tilde{X}(\infty))| \\
 &= |\mathbb{E}G_{\tilde{X}} f_h(\tilde{X}(\infty)) - \mathbb{E}G_Y f_h(\tilde{X}(\infty))| \\
 (3.9) \quad &\leq |\mathbb{E}G_{\tilde{X}} f_h(\tilde{X}(\infty)) - G_Y f_h(\tilde{X}(\infty))|.
 \end{aligned}$$

While the two random variables on the left side of (3.9) are usually defined on different probability spaces, the two random variables on the right side of (3.9) are both functions of $\tilde{X}(\infty)$. Thus, we have achieved a coupling through Lemma 1. Setting up the Poisson equation is a generic first step one performs any time one wishes to apply Stein's method to a problem. The next step is to bound the equivalent of our $|\mathbb{E}G_Y f_h(\tilde{X}(\infty))|$. This is usually done by using a coupling argument. However, this coupling is always problem specific, and is one of the greatest sources of difficulty one encounters when applying Stein's method. In our case, this generator coupling is natural because we deal with Markov processes $\tilde{X}$ and Y .

3.3. Taylor Expansion. To bound the right side of (3.9), we study the difference $G_{\tilde{X}} f_h(x) - G_Y f_h(x)$. For that we perform Taylor expansion on $G_{\tilde{X}} f_h(x)$. To illustrate this, suppose that $f_h''(x)$ exists for all $x \in \mathbb{R}$, and is absolutely continuous. Then for any $k \in \mathbb{Z}_+$, and $x = x_k = \delta(k - x(\infty))$, we recall that $b(x) = \delta(\lambda - d(k))$ in (3.8) to see that

$$\begin{aligned}
 G_{\tilde{X}} f_h(x) &= \lambda(f_h(x + \delta) - f_h(x)) + d(k)(f_h(x - \delta) - f_h(x)) \\
 &= f_h'(x)\delta(\lambda - d(k)) + \frac{1}{2}\delta^2 f_h''(x)(\lambda + d(k)) \\
 &\quad + \frac{1}{2}\lambda\delta^2(f_h''(\xi) - f_h''(x)) + \frac{1}{2}d(k)\delta^2(f_h''(\eta) - f_h''(x)) \\
 &= f_h'(x)b(x) + \frac{1}{2}\delta^2(2\lambda - \frac{1}{\delta}b(x))f_h''(x) \\
 &\quad + \frac{1}{2}\mu(f_h''(\xi) - f_h''(x)) + \frac{1}{2}(\lambda - \frac{1}{\delta}b(x))\delta^2(f_h''(\eta) - f_h''(x)) \\
 &= G_Y f_h(x) - \frac{1}{2}\delta f_h''(x)b(x) + \frac{1}{2}\mu(f_h''(\xi) - f_h''(x)) + \frac{1}{2}(\mu - \delta b(x))(f_h''(\eta) - f_h''(x)),
 \end{aligned}$$

where $\xi \in [x, x + \delta]$ and $\eta \in [x - \delta, x]$. We invoke the absolute continuity of $f_h''(x)$ to get

$$\begin{aligned}
 &|\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))| \\
 &\leq \frac{1}{2}\delta\mathbb{E}[|f_h''(\tilde{X}(\infty))b(\tilde{X}(\infty))|] + \frac{\mu}{2}\mathbb{E}\left[\int_{\tilde{X}(\infty)}^{\tilde{X}(\infty)+\delta} |f_h'''(y)| dy\right] \\
 (3.10) \quad &+ \frac{\mu}{2}\mathbb{E}\left[\int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f_h'''(y)| dy\right] + \frac{1}{2}\delta\mathbb{E}[|b(\tilde{X}(\infty))| \int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f_h'''(y)| dy].
 \end{aligned}$$

As one can see, to show that the right hand side of (3.10) vanishes as $\delta \rightarrow 0$, we must be able to bound the derivatives of $f_h(x)$; we refer to these as gradient bounds. Furthermore, we will also need bounds on moments of $|\tilde{X}(\infty)|$; we refer to these as moment bounds. Both moment and gradient bounds will vary between the Erlang-A or Erlang-C setting, and the gradient bounds will be different for the Wasserstein, and Kolmogorov settings. Moment bounds will be presented in Section 3.4, and gradient bounds in the Wasserstein setting will be presented in Section 3.5. We discuss the Kolmogorov setting separately in Section 5. In that case we face an added difficulty because $f_h''(x)$ has a discontinuity, meaning that we cannot use (3.10) directly.

3.4. Moment Bounds. We now state several lemmas that establish necessary moment bounds for showing that (3.10) is small. These moment bounds are used in the proofs of both the Wasserstein metric and the Kolmogorov metric versions of the theorems.

LEMMA 2. *Consider the Erlang-C model ($\alpha = 0$). For all $n \geq 1, \lambda > 0$, and $\mu > 0$ satisfying $0 < R < n$,*

$$(3.11) \quad \mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)] \leq \frac{4}{3} + \frac{2\delta^2}{3},$$

$$(3.12) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)|] \leq \sqrt{\frac{4}{3} + \frac{2\delta^2}{3}},$$

$$(3.13) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)|] \leq 2|\zeta|$$

$$(3.14) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)|] \leq \frac{1}{|\zeta|} + \frac{\delta^2}{4|\zeta|} + \frac{\delta}{2},$$

$$(3.15) \quad \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) \leq (2 + \delta)|\zeta|.$$

The proof of Lemma 2 is provided in Section B.1. Note that the event $\{\tilde{X}(\infty) \leq -\zeta\}$ is equal to the event $\{X(\infty) \leq n\}$. In particular, (3.15) gives a bound on the probability that no customers are waiting in the system.

Observe that we have the freedom to use either (3.12) or (3.13), depending on which bound suits our needs better in any given situation. We see that (3.14) immediately implies that when $\delta \leq 1$,

$$\begin{aligned} |\zeta| \mathbb{P}(\tilde{X}(\infty) \geq -\zeta) &\leq |\zeta| \wedge \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)|] \\ &\leq |\zeta| \wedge \left(\frac{1}{|\zeta|} + \frac{\delta^2}{4|\zeta|} + \frac{\delta}{2} \right) \\ (3.16) \quad &\leq 7/4, \end{aligned}$$

where to get the last inequality we considered separately the cases where $|\zeta| \leq 1$ and $|\zeta| \geq 1$. This bound will be used in the proofs of Theorems 1 and 3. The following lemma states

the necessary moment bounds for the Erlang-A model. The underloaded and overloaded cases have to be handled separately. Since the drift $b(x)$ is different between the Erlang-A and Erlang-C models, the quantities bounded in the following lemma will resemble those in Lemma 2, but will not be identical. A proof outline for Lemma 3 will be provided in Section B.2 for the underloaded case, and in Section B.3 for the overloaded case.

LEMMA 3. *Consider the Erlang-A model ($\alpha > 0$). Fix $n \geq 1, \lambda > 0, \mu > 0$, and $\alpha > 0$. If $0 < R \leq n$ (an underloaded system), then*

$$(3.17) \quad \mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)] \leq \frac{1}{3} \left(\frac{\alpha}{\mu} \delta^2 + \delta^2 + 4 \right),$$

$$(3.18) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)|] \leq \sqrt{\frac{1}{3} \left(\frac{\alpha}{\mu} \delta^2 + \delta^2 + 4 \right)},$$

$$(3.19) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)|] \leq 2|\zeta| + \frac{\alpha}{\mu} \sqrt{\frac{1}{3} \left(\frac{\mu}{\alpha} \delta^2 + \frac{\mu}{\alpha} 4 + \delta^2 \right)},$$

$$(3.20) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)|] \leq \left(1 + \frac{\delta^2}{4} + \frac{\delta}{2} \sqrt{\frac{1}{3} \left(\frac{\alpha}{\mu} \delta^2 + \delta^2 + 4 \right)} \right) \left(\frac{\mu}{\mu \wedge \alpha} \wedge \frac{1}{|\zeta|} \right),$$

$$(3.21) \quad \mathbb{E}[(\tilde{X}(\infty) + \zeta)^2 1(\tilde{X}(\infty) \geq -\zeta)] \leq \frac{1}{3} \left(\frac{\mu}{\alpha} \delta^2 + \frac{\mu}{\alpha} 4 + \delta^2 \right),$$

$$(3.22) \quad \mathbb{E}[(\tilde{X}(\infty) + \zeta) 1(\tilde{X}(\infty) \geq -\zeta)] \leq \sqrt{\frac{1}{3} \left(\frac{\mu}{\alpha} \delta^2 + \frac{\mu}{\alpha} 4 + \delta^2 \right)},$$

$$(3.23) \quad \mathbb{E}[(\tilde{X}(\infty) + \zeta) 1(\tilde{X}(\infty) \leq -\zeta)] \leq \frac{1}{|\zeta|} \left(\frac{\delta^2}{4} \frac{\alpha}{\mu} + \frac{\delta^2}{4} + 1 \right),$$

$$(3.24) \quad \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) \leq (2 + \delta) \left(|\zeta| + \frac{\alpha}{\mu} \sqrt{\frac{1}{3} \left(\frac{\mu}{\alpha} \delta^2 + \frac{\mu}{\alpha} 4 + \delta^2 \right)} \right).$$

and if $n \leq R$ (an overloaded system), then

$$(3.25) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)|] \leq \sqrt{\frac{1}{\alpha \wedge \mu} \left(\alpha \frac{\delta^2}{4} + \mu \right)},$$

$$(3.26) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)|] \leq \frac{1}{|\zeta|} \left(\frac{\delta^2}{4} + \frac{\mu}{\alpha} \right),$$

$$(3.27) \quad \mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \geq -\zeta)] \leq \frac{1}{3} \left(\delta^2 + 4 \frac{\mu}{\alpha} \right),$$

$$(3.28) \quad \mathbb{E}[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)|] \leq \sqrt{\frac{1}{3} \left(\delta^2 + 4 \frac{\mu}{\alpha} \right)},$$

$$(3.29) \quad \mathbb{E}[|(\tilde{X}(\infty) + \zeta) 1(\tilde{X}(\infty) \leq -\zeta)|] \leq \frac{1}{|\zeta|} \left(\frac{\delta^2}{4} + 1 \right),$$

$$(3.30) \quad \mathbb{E}[(\tilde{X}(\infty) + \zeta)^2 1(\tilde{X}(\infty) \leq -\zeta)] \leq \frac{\delta^2}{4} \frac{\alpha}{\mu} + 1,$$

$$(3.31) \quad \mathbb{E} \left[|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) \leq -\zeta)| \right] \leq \sqrt{\frac{\delta^2}{4} \frac{\alpha}{\mu} + 1},$$

$$(3.32) \quad \mathbb{E} \left[|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) \leq -\zeta)| \right] \leq \frac{\alpha}{\mu} \sqrt{\frac{1}{3} \left(\delta^2 + 4 \frac{\mu}{\alpha} \right)},$$

$$(3.33) \quad \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) \leq (3 + \delta) \frac{16}{\sqrt{2}} \left(\frac{\delta^2}{4} + 1 \right) \left(\left(\frac{1}{\zeta} \vee \frac{\alpha}{\mu} \right) \wedge \sqrt{\frac{\alpha}{\mu}} \right).$$

3.5. *Wasserstein Gradient Bounds.* Recall the Poisson equation (3.3). In this section, we present the gradient bounds on the solutions to the Poisson equation in the Wasserstein setting ($\mathcal{H} = \text{Lip}(1)$). In Section 4 we prove Theorems 1 and 2 by applying these gradient bounds together with the moment bounds from Section 3.4 to (3.10).

The following lemmas present the Wasserstein gradient bounds. We recall that $R \leq n$ implies $\zeta \leq 0$, and $R \geq n$ implies $\zeta \geq 0$. For both lemmas, we recall the Poisson equation (3.3) and that the family of solutions to this equation is given by (3.4); in particular, this family is parametrized by constants $a_1, a_2 \in \mathbb{R}$. We fix $h(x) \in \mathcal{H} = \text{Lip}(1)$, and let $f_h(x)$ be a solution to the Poisson equation.

LEMMA 4. *Consider the Erlang-C model ($\alpha = 0$). The solution to the Poisson equation $f_h(x)$ is twice continuously differentiable, with an absolutely continuous second derivative. Fix a solution in (3.4) with parameter $a_2 = 0$. Then for all $n \geq 1, \lambda > 0$, and $\mu > 0$ satisfying $0 < R < n$,*

$$(3.34) \quad |f'_h(x)| \leq \begin{cases} \frac{1}{\mu} (6.5 + 4.2/|\zeta|), & x \leq -\zeta, \\ \frac{1}{\mu} \frac{1}{|\zeta|} (x + 1 + 2/|\zeta|), & x \geq -\zeta. \end{cases}$$

$$(3.35) \quad |f''_h(x)| \leq \begin{cases} \frac{32}{\mu} (1 + 1/|\zeta|), & x \leq -\zeta, \\ \frac{1}{\mu|\zeta|}, & x \geq -\zeta, \end{cases}$$

and for those $x \in \mathbb{R}$ where $f'''_h(x)$ exists,

$$(3.36) \quad |f'''_h(x)| \leq \begin{cases} \frac{1}{\mu} (23 + 13/|\zeta|), & x \leq -\zeta, \\ 2/\mu, & x \geq -\zeta. \end{cases}$$

LEMMA 5. *Consider the Erlang-A model ($\alpha > 0$). The solution to the Poisson equation $f_h(x)$ is twice continuously differentiable, with an absolutely continuous second derivative. Fix a solution in (3.4) with parameter $a_2 = 0$. Then there exists a constant $C > 0$ independent of λ, n, μ , and α such that for all $n \geq 1, \lambda > 0, \mu > 0$, and $\alpha > 0$ satisfying $0 < R \leq n$*

(an underloaded system),

$$(3.37) \quad |f'_h(x)| \leq \begin{cases} C \left(\sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} + 1 \right) \frac{1}{\mu}, & x \leq -\zeta, \\ C \left(\frac{\mu}{\alpha} + \sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} + 1 \right) \frac{1}{\mu}, & x \geq -\zeta, \end{cases}$$

$$(3.38) \quad |f''_h(x)| \leq \begin{cases} C \left(\sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} + 1 \right) \frac{1}{\mu}, & x \leq 0, \\ C \left[\left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \left(\sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} + 1 \right) + 1 \right] \frac{1}{\mu}, & x \in [0, -\zeta], \\ C \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \left(\sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} \right) \frac{1}{\mu}, & x \geq -\zeta, \end{cases}$$

(3.39)

and for those $x \in \mathbb{R}$ where $f'''_h(x)$ exists,

$$(3.40) \quad |f'''_h(x)| \leq \begin{cases} C \left(\sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} + 1 \right) \frac{1}{\mu}, & x \leq 0, \\ C \left(\sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} + \frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \frac{1}{\mu}, & x \in [0, -\zeta], \\ C \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \frac{1}{\mu}, & x \geq -\zeta, \end{cases}$$

and for all $n \geq 1, \lambda > 0, \mu > 0$, and $\alpha > 0$ satisfying $n \leq R$ (an overloaded system),

$$(3.41) \quad |f'_h(x)| \leq \begin{cases} C \left(\frac{1}{\mu} + \frac{1}{\sqrt{\alpha}} \frac{1}{\sqrt{\mu}} + \frac{\zeta}{\mu} \wedge \frac{1}{\alpha} \right), & x \leq -\zeta, \\ C \left(\frac{1}{\mu} + \frac{1}{\sqrt{\alpha}} \frac{1}{\sqrt{\mu}} + \frac{1}{\alpha} \right), & x \geq -\zeta, \end{cases}$$

$$(3.42) \quad |f''_h(x)| \leq \begin{cases} C \left(\frac{1}{\mu} + \frac{1}{\sqrt{\alpha}} \frac{1}{\sqrt{\mu}} + \frac{\zeta}{\mu} \wedge \frac{1}{\alpha} \right), & x \leq -\zeta, \\ C \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \frac{1}{\mu} |x| + C \left(\frac{1}{\mu} + \frac{1}{\sqrt{\alpha}} \frac{1}{\sqrt{\mu}} \right), & x \geq -\zeta, \end{cases}$$

(3.43)

and for those $x \in \mathbb{R}$ where $f'''_h(x)$ exists,

$$(3.44) \quad |f'''_h(x)| \leq \frac{C}{\mu} \left(1 + \sqrt{\frac{\mu}{\alpha}} + \zeta \wedge \frac{\mu}{\alpha} \right), \quad x \leq -\zeta,$$

$$(3.45) \quad |f'''_h(x)| \leq \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \left(1 + \frac{\alpha}{\mu} x^2 \right) + \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} \right) |x|, \quad x \geq -\zeta,$$

$$(3.46) \quad |f'''_h(x)| \leq \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) + \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right)^2 |x|, \quad x \geq -\zeta.$$

REMARK 3. These lemmas validate the Taylor expansion used to obtain (3.10) in the Wasserstein setting, because $f''_h(x)$ is absolutely continuous. Furthermore, the functions $f_h(x)$ considered in these lemmas all satisfy the conditions of Lemma 1, because $f'_h(x)$ grows at most linearly in both lemmas.

Lemma 4 is proved in Section C.1. The procedure to prove Lemma 4 is very similar to the one used for Lemma 5, and we do not provide a full proof for the latter lemma. Instead, we outline the necessary steps needed for a proof, and emphasize all the differences with the proof of Lemma 4; this is done in Section C.1.2 for the underloaded system and Section C.1.3 for the overloaded system. Since a full proof is not provided, we do not present explicit constants in the gradient bounds of Lemma 5, although they can certainly be recovered.

In establishing these gradient bounds, we do not rely on the Lyapunov function methods from [9]. Instead, we analyze the closed form solution of the Poisson equation directly; however, this is only possible because the Poisson equation is an ordinary differential equation. Typically, this direct analysis cannot be generalized to when the Poisson equation is a partial differential equation, i.e. when the diffusion process Y is multidimensional. We are now ready to prove Theorems 1 and 2.

4. Proof of Theorems 1 and 2: the Wasserstein metric. In this section we prove Theorems 1 and 2. Fix $h(x) \in \text{Lip}(1)$, and recall that the family of solutions to the Poisson equation is given by (3.4). For the remainder of Section 4, we fix one such solution $f_h(x)$ with $a_2 = 0$. Then by Lemmas 4 and 5, $f_h''(x)$ is absolutely continuous, implying that (3.10) holds; we recall it here as

$$\begin{aligned}
 & \left| \mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty)) \right| \\
 & \leq \frac{1}{2}\delta \mathbb{E} \left[|f_h''(\tilde{X}(\infty))b(\tilde{X}(\infty))| \right] + \frac{\mu}{2} \mathbb{E} \left[\int_{\tilde{X}(\infty)}^{\tilde{X}(\infty)+\delta} |f_h'''(y)| dy \right] \\
 (4.1) \quad & + \frac{\mu}{2} \mathbb{E} \left[\int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f_h'''(y)| dy \right] + \frac{1}{2}\delta \mathbb{E} \left[|b(\tilde{X}(\infty))| \int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f_h'''(y)| dy \right].
 \end{aligned}$$

The proof of Theorems 1 and 2 simply involves applying the moment bounds and gradient bounds to show that the error bound in (4.1) is small.

PROOF OF THEOREM 1. Throughout the proof we assume that $R \geq 1$, or equivalently, $\delta \leq 1$. We bound each of the terms on the right side of (4.1) individually. We recall here that the support of $\tilde{X}(\infty)$ is a δ -spaced grid, and in particular this grid contains the point $-\zeta$. In the bounds that follow, we will often consider separately the cases where $\tilde{X}(\infty) \leq -\zeta - \delta$, and $\tilde{X}(\infty) \geq -\zeta$. We recall that

$$(4.2) \quad b(x) = \mu[(x + \zeta)^- - \zeta^-] = \begin{cases} -\mu x, & x \leq -\zeta, \\ \mu \zeta, & x \geq -\zeta, \end{cases}$$

and apply the moment bounds (3.12), (3.13), and the gradient bound (3.35), to see that

$$\begin{aligned}
\mathbb{E}\left[|f_h''(\tilde{X}(\infty))b(\tilde{X}(\infty))|\right] &\leq 32(1 + 1/|\zeta|)\mathbb{E}\left[|\tilde{X}(\infty)|1(\tilde{X}(\infty) \leq -\zeta - \delta)\right] + \mathbb{P}(\tilde{X}(\infty) \geq -\zeta) \\
&\leq 32(1 + 1/|\zeta|)\left(2|\zeta| \wedge \sqrt{\frac{4}{3} + \frac{2\delta^2}{3}}\right) + 1 \\
&\leq 32\left(\sqrt{\frac{4}{3} + \frac{2\delta^2}{3}} + 2\right) + 1 \\
&\leq 32(\sqrt{2} + 2) + 1 \leq 111.
\end{aligned}$$

Next, we use (3.15) and the gradient bound in (3.36) to get

$$\begin{aligned}
\frac{\mu}{2}\mathbb{E}\left[\int_{\tilde{X}(\infty)}^{\tilde{X}(\infty)+\delta}|f_h'''(y)|dy\right] &\leq \frac{\delta}{2}\left((23 + 13/|\zeta|)\mathbb{P}(\tilde{X}(\infty) \leq -\zeta - \delta) + 2\mathbb{P}(\tilde{X}(\infty) \geq -\zeta)\right) \\
&\leq \frac{\delta}{2}\left(23 + \frac{13}{|\zeta|}(3|\zeta|)\right) \leq 31\delta.
\end{aligned}$$

By a similar argument, we can show that

$$\frac{\mu}{2}\mathbb{E}\left[\int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)}|f_h'''(y)|dy\right] \leq 31\delta,$$

with the only difference in the argument being that we consider the cases when $\tilde{X}(\infty) \leq -\zeta$ and $\tilde{X}(\infty) \geq -\zeta + \delta$, instead of $\tilde{X}(\infty) \leq -\zeta - \delta$ and $\tilde{X}(\infty) \geq -\zeta$. Lastly, we use the form of $b(x)$, the moment bounds (3.12), (3.13), and (3.16), and the gradient bound (3.36) to get

$$\begin{aligned}
&\frac{\delta}{2}\mathbb{E}\left[|b(\tilde{X}(\infty))|\int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)}|f_h'''(y)|dy\right] \\
&\leq \frac{\delta^2}{2}\left((23 + 13/|\zeta|)\mathbb{E}\left[|\tilde{X}(\infty)|1(\tilde{X}(\infty) \leq -\zeta)\right] + 2|\zeta|\mathbb{P}(\tilde{X}(\infty) \geq -\zeta + \delta)\right) \\
&\leq \frac{\delta^2}{2}\left((23 + 13/|\zeta|)\left(2|\zeta| \wedge \sqrt{\frac{4}{3} + \frac{2\delta^2}{3}}\right) + 14/4\right) \\
&\leq \frac{\delta^2}{2}\left(23\sqrt{2} + 26 + 14/4\right) \leq 32\delta^2.
\end{aligned}$$

Hence, from (3.10) we conclude that for all $R \geq 1$, and $h(x) \in \text{Lip}(1)$,

$$(4.3) \quad \left|\mathbb{E}h(\tilde{X}(\infty)) - \mathbb{E}h(Y(\infty))\right| \leq \delta(111 + 31 + 31 + 32\delta) \leq 205\delta,$$

which proves Theorem 1. □

Proving Theorem 2 consists of bounding the four error terms in (3.10). Since the procedure is very similar to the proof of Theorem 1, we will only outline which gradient and moment bounds need to be used to bound each error term.

We start with the underloaded case, when $R \leq n$. To bound the first term in (3.10), we use moment bounds (3.18), (3.19), and (3.22), together with the gradient bounds in (3.38). For the second and third terms, we use moment bound (3.24) and the gradient bounds in (3.40). For the fourth term, we use moment bounds (3.18)–(3.22), and the gradient bounds in (3.40).

We now prove the overloaded case, when $R \geq n$. To bound the first term in (3.10), we use moment bounds (3.25)–(3.32), together with the gradient bounds in (3.42). For the second and third terms, we use moment bounds (3.27), (3.28), and (3.33), together with the gradient bounds in (3.44) and (3.45). For the fourth term, we use moment bounds (3.25)–(3.32), and gradient bounds in (3.44) and (3.46).

5. The Kolmogorov Metric. In this section we prove Theorems 3 and 4, which are stated in the Kolmogorov setting. The biggest difference between the Wasserstein and Kolmogorov settings is that in the latter, the test functions $h(x)$ used in the Poisson equation (3.3) are discontinuous. For this reason, the gradient bounds from Lemmas 4 and 5 do not hold anymore, and new gradient bounds need to be derived separately for the Kolmogorov setting; we present these new gradient bounds in Section 5.1. Furthermore, the solution to the Poisson equation no longer has a continuous second derivative, meaning that the Taylor expansion we used to derive the upper bound in (3.10) is invalid. We discuss an alternative to (3.10) in Section 5.2. This alternative bound contains a new error term that cannot be handled by the gradient bounds, nor the moment bounds. This term appears because the solution to the Poisson equation has a discontinuous second derivative, and to bound it we present Lemma 8. We then prove Theorems 3 and 4 in Section 5.3.

5.1. Kolmogorov Gradient Bounds. Recall that in the Kolmogorov setting, we take the class of test functions for the Poisson equation (3.3) to be $\mathcal{H}_K$ defined in (1.8). For the statement of the following two lemmas, we fix $a \in \mathbb{R}$ and set $h(x) = 1_{(-\infty, a]}(x)$. We use $f_a(x)$ instead of $f_h(x)$ to denote a solution to the Poisson equation. We recall that the family of solutions to the Poisson equation is parametrized by constants $a_1, a_2 \in \mathbb{R}$. The following lemmas state the gradient bounds in the Kolmogorov setting.

LEMMA 6. *Consider the Erlang-C model ($\alpha = 0$). Any solution to the Poisson equation $f_a(x)$ is continuously differentiable, with an absolutely continuous derivative. Fix a solution in (3.4) with parameter $a_2 = 0$. Then for all $n \geq 1, \lambda > 0$, and $\mu > 0$ satisfying $0 < R < n$,*

$$(5.1) \quad |f'_a(x)| \leq \begin{cases} 5/\mu, & x \leq -\zeta, \\ \frac{1}{\mu|\zeta|}, & x \geq -\zeta, \end{cases}$$

and for all $x \in \mathbb{R}$,

$$(5.2) \quad |f_a''(x)| \leq 3/\mu,$$

where $f_a''(x)$ is understood to be the left derivative at the point $x = a$.

LEMMA 7. Consider the Erlang-A model ($\alpha > 0$). Any solution to the Poisson equation $f_a(x)$ is continuously differentiable, with an absolutely continuous derivative. Fix a solution in (3.4) with parameter $a_2 = 0$, and fix $n \geq 1, \lambda > 0, \mu > 0$, and $\alpha > 0$. If $0 < R \leq n$ (an underloaded system), then

$$(5.3) \quad |f_a'(x)| \leq \begin{cases} \frac{1}{\mu} \sqrt{2\pi} e^{1/2}, & x \leq -\zeta, \\ \frac{1}{\mu} \left(\sqrt{\frac{\pi}{2}} \frac{\mu}{\alpha} \wedge \frac{1}{|\zeta|} \right), & x \geq -\zeta, \end{cases}$$

and if $n \leq R$ (an overloaded system), then

$$(5.4) \quad |f_a'(x)| \leq \begin{cases} \frac{1}{\mu} \sqrt{\frac{\pi}{2}}, & x \leq -\zeta, \\ \frac{1}{\mu} \sqrt{\frac{\pi}{2}} \left(1 + \sqrt{\frac{\mu}{\alpha}} \right), & x \geq -\zeta. \end{cases}$$

Moreover, for all $\lambda > 0, n \geq 1, \mu > 0$, and $\alpha > 0$, and all $x \in \mathbb{R}$,

$$(5.5) \quad |f_a''(x)| \leq 3/\mu,$$

where $f_a''(x)$ is understood to be the left derivative at the point $x = a$.

Lemmas 6 and 7 are proved in Section C.2. Unlike the Wasserstein setting, these lemmas do not guarantee that $f_a''(x)$ is absolutely continuous. Indeed, for any $a \in \mathbb{R}$, we can rewrite the Poisson equation as

$$\mu f_a''(x) = \mathbb{P}(Y(\infty) \leq a) - 1_{(-\infty, a]}(x) - b(x) f_a'(x),$$

and see that $f_a''(x)$ is in fact discontinuous at the point $x = a$. This means that we cannot directly use the error bound in (3.10), and require a different expansion of $G_{\tilde{X}} f_a(x)$.

5.2. *Alternative Taylor Expansion.* To get an error bound similar to (3.10), we first define

$$(5.6) \quad \epsilon_1(x) = \int_x^{x+\delta} (x + \delta - y)(f_a''(y) - f_a''(x-)) dy,$$

$$(5.7) \quad \epsilon_2(x) = \int_{x-\delta}^x (y - (x - \delta))(f_a''(y) - f_a''(x-)) dy.$$

Now observe that

$$\begin{aligned}
f_a(x + \delta) - f_a(x) &= f'_a(x)\delta + \int_x^{x+\delta} (x + \delta - y)f''_a(y)dy \\
&= f'_a(x)\delta + \frac{1}{2}\delta^2 f''_a(x-) + \int_x^{x+\delta} (x + \delta - y)(f''_a(y) - f''_a(x-))dy \\
&= f'_a(x)\delta + \frac{1}{2}\delta^2 f''_a(x-) + \epsilon_1(x),
\end{aligned}$$

and

$$\begin{aligned}
(f_a(x - \delta) - f_a(x)) &= -f'_a(x)\delta + \int_{x-\delta}^x (y - (x - \delta))f''_a(y)dy \\
&= -f'_a(x)\delta + \frac{1}{2}\delta^2 f''_a(x-) + \int_{x-\delta}^x (y - (x - \delta))(f''_a(y) - f''_a(x-))dy \\
&= -f'_a(x)\delta + \frac{1}{2}\delta^2 f''_a(x-) + \epsilon_2(x).
\end{aligned}$$

For $k \in \mathbb{Z}_+$ and $x = x_k = \delta(k - x(\infty))$, we recall the forms of $G_Y f_a(x)$ and $G_{\tilde{X}} f_a(x)$ from (3.2) and (3.6) to see that

$$\begin{aligned}
G_{\tilde{X}} f_a(x) &= \lambda \delta f'_a(x) + \lambda \frac{1}{2} \delta^2 f''_a(x-) + \lambda \epsilon_1(x) \\
&\quad - d(k) \delta f'_a(x) + d(k) \frac{1}{2} \delta^2 f''_a(x-) + d(k) \epsilon_2(x) \\
&= b(x) f'_a(x) + \lambda \frac{1}{2} \delta^2 f''_a(x-) + \lambda \epsilon_1(x) \\
&\quad + \left(\lambda - \frac{1}{\delta} b(x)\right) \frac{1}{2} \delta^2 f''_a(x-) + \left(\lambda - \frac{1}{\delta} b(x)\right) \epsilon_2(x) \\
&= G_Y f(x) - b(x) \frac{1}{2} \delta f''_a(x-) + \lambda(\epsilon_1(x) + \epsilon_2(x)) - \frac{1}{\delta} b(x) \epsilon_2(x),
\end{aligned}$$

where in the second equality we used the fact that $b(x) = \delta(\lambda - d(k))$, and in the last equality we use that $\delta^2 \lambda = \mu$. Combining this with (3.9), we have an error bound similar to (3.10):

$$\begin{aligned}
&\left| \mathbb{P}(\tilde{X}(\infty) \leq a) - \mathbb{P}(Y(\infty) \leq a) \right| \\
&\leq \frac{1}{2} \delta \mathbb{E} \left[\left| f''_a(\tilde{X}(\infty)-) b(\tilde{X}(\infty)) \right| \right] + \lambda \mathbb{E} \left[\left| \epsilon_1(\tilde{X}(\infty)) \right| \right] \\
(5.8) \quad &\quad + \lambda \mathbb{E} \left[\left| \epsilon_2(\tilde{X}(\infty)) \right| \right] + \frac{1}{\delta} \mathbb{E} \left[\left| b(\tilde{X}(\infty)) \epsilon_2(\tilde{X}(\infty)) \right| \right],
\end{aligned}$$

where $\epsilon_1(x)$ and $\epsilon_2(x)$ are as in (5.6) and (5.7). To bound the error terms in (5.8) that are associated with $\epsilon_1(x)$ and $\epsilon_2(x)$, we need to analyze the difference $f''_a(y) - f''_a(x-)$ for

$|x - y| \leq \delta$. Since $f_a(x)$ is a solution to the Poisson equation (3.3), we see that for any $x, y \in \mathbb{R}$ with $y \neq a$,

$$f_a''(y) - f_a''(x-) = \frac{1}{\mu} [1_{(-\infty, a]}(x) - 1_{(-\infty, a]}(y) + b(x)f_a'(x) - b(y)f_a'(y)].$$

Therefore, for any $y \in [x, x + \delta]$ with $y \neq a$,

$$\begin{aligned} |f_a''(y) - f_a''(x-)| &\leq \frac{1}{\mu} [1_{(a-\delta, a]}(x) + |b(x)| |f_a'(x) - f_a'(y)| + |b(x) - b(y)| |f_a'(y)|] \\ (5.9) \quad &\leq \frac{1}{\mu} [1_{(a-\delta, a]}(x) + \delta |b(x)| \|f''\| + |b(x) - b(y)| |f_a'(y)|], \end{aligned}$$

and likewise, for any $y \in [x - \delta, x]$ with $y \neq a$,

$$\begin{aligned} |f_a''(y) - f_a''(x-)| &\leq \frac{1}{\mu} [1_{(a, a+\delta]}(x) + |b(x)| |f_a'(x) - f_a'(y)| + |b(x) - b(y)| |f_a'(y)|] \\ (5.10) \quad &\leq \frac{1}{\mu} [1_{(a, a+\delta]}(x) + \delta |b(x)| \|f''\| + |b(x) - b(y)| |f_a'(y)|]. \end{aligned}$$

Note that the inequalities above contain the indicators $1_{(a-\delta, a]}(x)$ and $1_{(a, a+\delta]}(x)$. When we consider the upper bound in (5.8), these indicators will manifest themselves as the probabilities $\mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a)$ and $\mathbb{P}(a < \tilde{X}(\infty) \leq a + \delta)$. To this end we present the following lemma, which will be used in the proof of Theorem 3.

LEMMA 8. *Consider the Erlang-C model ($\alpha = 0$). Let W be an arbitrary random variable with cumulative distribution function $F_W : \mathbb{R} \rightarrow [0, 1]$. Let $\omega(F_W)$ be the modulus of continuity of F_W , defined as*

$$\omega(F_W) = \sup_{\substack{x, y \in \mathbb{R} \\ x \neq y}} \frac{|F_W(x) - F_W(y)|}{|x - y|}.$$

Recall that $d_K(\tilde{X}(\infty), W)$ is the Kolmogorov distance between $X(\infty)$ and W . Then for any $a \in \mathbb{R}$, $n \geq 1$, and $0 < R < n$,

$$\mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a + \delta) \leq \omega(F_W)2\delta + d_K(\tilde{X}(\infty), W) + 9\delta^2 + 8\delta^4.$$

This lemma is proved in Section D.1. We will apply Lemma 8 with $W = Y(\infty)$ in the proof of Theorem 3 that follows. The following lemma guarantees that the modulus of continuity of the cumulative distribution function of $Y(\infty)$ is bounded by a constant independent of λ, n , and μ . Its proof is provided in Section D.2.

LEMMA 9. *Consider the Erlang-C model ($\alpha = 0$), and let $\nu(x)$ be the density of $Y(\infty)$. Then for all $n \geq 1, \lambda > 0$, and $\mu > 0$ satisfying $0 < R < n$,*

$$|\nu(x)| \leq \sqrt{\frac{2}{\pi}}, \quad x \in \mathbb{R}.$$

Lemmas 8 and 9 are stated for the Erlang-C model, but one can easily repeat the arguments in the proofs of those lemmas to prove analogues for the Erlang-A model. Therefore, we state the following lemmas without proof.

LEMMA 10. *Consider the Erlang-A model ($\alpha > 0$). Let W be an arbitrary random variable with cumulative distribution function $F_W : \mathbb{R} \rightarrow [0, 1]$. Let $\omega(F_W)$ be the modulus of continuity of F_W . Then for any $a \in \mathbb{R}$, $\alpha > 0$, $n \geq 1$, and $R > 0$,*

$$\mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a + \delta) \leq \omega(F_W)2\delta + d_K(\tilde{X}(\infty), W) + 9\left(\frac{\alpha}{\mu} \vee 1\right)\delta^2 + 8\left(\frac{\alpha}{\mu} \vee 1\right)^2\delta^4.$$

LEMMA 11. *Consider the Erlang-A model ($\alpha > 0$), and let $\nu(x)$ be the density of $Y(\infty)$. Fix $n \geq 1$, $\lambda > 0$, $\mu > 0$, and $\alpha > 0$. If $0 < R \leq n$, then*

$$|\nu(x)| \leq \sqrt{\frac{2}{\pi}}, \quad x \in \mathbb{R},$$

and if $n \leq R$, then

$$|\nu(x)| \leq \sqrt{\frac{2}{\pi}} \sqrt{\frac{\alpha}{\mu}}, \quad x \in \mathbb{R}.$$

5.3. Proof of Theorems 3 and 4.

PROOF OF THEOREM 3. Throughout the proof we assume that $R \geq 1$, or equivalently, $\delta \leq 1$. For $h(x) = 1_{(-\infty, a]}(x)$, we let $f_a(x)$ be a solution the Poisson equation (3.3) with parameter $a_2 = 0$. In this proof we will show that for all $a \in \mathbb{R}$,

$$(5.11) \quad \left| \mathbb{P}(\tilde{X}(\infty) \leq a) - \mathbb{P}(Y(\infty) \leq a) \right| \leq \frac{1}{2} \mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a + \delta) + 75\delta,$$

The upper bound in (5.11) is similar to (4.3), however (5.11) has the extra term

$$\frac{1}{2} \mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a + \delta).$$

The reason this term appears in the Kolmogorov setting but not in the Wasserstein setting is because $f_a''(x)$ is discontinuous in the Kolmogorov case, as opposed to the Wasserstein case where $f_h''(x)$ is continuous. Applying Lemmas 8 and 9 to the right hand side of (5.11), and taking the supremum over all $a \in \mathbb{R}$ on both sides, we see that

$$d_K(\tilde{X}(\infty), Y(\infty)) \leq \frac{1}{2} d_K(\tilde{X}(\infty), Y(\infty)) + 2\sqrt{\frac{2}{\pi}}\delta + 9\delta^2 + 8\delta^4 + 75\delta,$$

or

$$d_K(\tilde{X}(\infty), Y(\infty)) \leq 188\delta.$$

To prove Theorem 3 it remains to verify (5.11), which we now do. The argument we will use is similar to the argument used to prove (4.3) in Theorem 1. We will bound each of the terms in (5.8), which we recall here as

$$\begin{aligned} & \left| \mathbb{P}(\tilde{X}(\infty) \leq a) - \mathbb{P}(Y(\infty) \leq a) \right| \\ & \leq \frac{1}{2} \delta \mathbb{E} \left[\left| f_a''(\tilde{X}(\infty)) - b(\tilde{X}(\infty)) \right| \right] + \lambda \mathbb{E} \left[\left| \epsilon_1(\tilde{X}(\infty)) \right| \right] \\ & \quad + \lambda \mathbb{E} \left[\left| \epsilon_2(\tilde{X}(\infty)) \right| \right] + \frac{1}{\delta} \mathbb{E} \left[\left| b(\tilde{X}(\infty)) \epsilon_2(\tilde{X}(\infty)) \right| \right]. \end{aligned}$$

We also recall the form of $b(x)$ from (4.2). We use the moment bounds (3.12) and (3.16), and the gradient bound (5.2) to see that

$$\begin{aligned} \mathbb{E} \left[\left| f_a''(\tilde{X}(\infty)) - b(\tilde{X}(\infty)) \right| \right] & \leq \frac{3}{\mu} \mathbb{E} \left[\left| b(\tilde{X}(\infty)) \right| \right] \\ & = 3 \mathbb{E} \left[\left| \tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta - \delta) \right| \right] + 3 |\zeta| \mathbb{P}(\tilde{X}(\infty) \geq -\zeta) \\ & \leq 3 \sqrt{\frac{4}{3} + \frac{2\delta^2}{3}} + 3 \left(|\zeta| \wedge \mathbb{E} \left[\left| \tilde{X}(\infty) \right| 1(\tilde{X}(\infty) \geq -\zeta) \right] \right) \\ (5.12) \quad & \leq 3\sqrt{2} + \frac{21}{4} \leq 10. \end{aligned}$$

Next, we use (5.9), (5.12), and the gradient bound (5.1) to get

$$\begin{aligned} \lambda \mathbb{E} \left[\left| \epsilon_1(\tilde{X}(\infty)) \right| \right] & = \lambda \mathbb{E} \left[\int_{\tilde{X}(\infty)}^{\tilde{X}(\infty)+\delta} (\tilde{X}(\infty) + \delta - y) \left| f_a''(y) - f_a''(\tilde{X}(\infty)) \right| dy \right] \\ & \leq \frac{\lambda}{\mu} \mathbb{E} \left[1_{(a-\delta, a]}(\tilde{X}(\infty)) \int_{\tilde{X}(\infty)}^{\tilde{X}(\infty)+\delta} (\tilde{X}(\infty) + \delta - y) dy \right] \\ & \quad + \frac{\lambda}{\mu} \delta^3 \mathbb{E} \left[\left| b(\tilde{X}(\infty)) \right| \right] \|f_a''\| + \frac{\lambda}{\mu} \delta \mathbb{E} \left[\int_{\tilde{X}(\infty)}^{\tilde{X}(\infty)+\delta} |b(\tilde{X}(\infty)) - b(y)| \left| f_a'(y) \right| dy \right] \\ & \leq \frac{1}{2} \mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a) + 10\delta + 5\delta \\ & \equiv \frac{1}{2} \mathbb{P}(a - \delta < \tilde{X}(\infty) \leq a) + 15\delta, \end{aligned}$$

where in the last inequality we used the fact that for $y \in [\tilde{X}(\infty), \tilde{X}(\infty) + \delta]$,

$$b(\tilde{X}(\infty)) - b(y) = \mu \delta 1(\tilde{X}(\infty) \leq -\zeta - \delta).$$

By a similar argument, one can check that

$$\lambda \mathbb{E} \left[\left| \epsilon_2(\tilde{X}(\infty)) \right| \right] \leq \frac{1}{2} \mathbb{P}(a < \tilde{X}(\infty) \leq a + \delta) + 15\delta,$$

with the only difference in the argument being that we consider the cases when $\tilde{X}(\infty) \leq -\zeta$ and $\tilde{X}(\infty) \geq -\zeta + \delta$, instead of $\tilde{X}(\infty) \leq -\zeta - \delta$ and $\tilde{X}(\infty) \geq -\zeta$. Lastly, we use the first inequality in (5.10) to see that

$$\begin{aligned}
& \frac{1}{\delta} \mathbb{E} \left[|b(\tilde{X}(\infty)) \epsilon_2(\tilde{X}(\infty))| \right] \\
& \leq \frac{1}{\mu} \mathbb{E} \left[|b(\tilde{X}(\infty))| \int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} \left[1_{(a, a+\delta]}(\tilde{X}(\infty)) + |b(\tilde{X}(\infty))| \left(|f'_a(\tilde{X}(\infty))| + |f'_a(y)| \right) \right. \right. \\
& \quad \left. \left. + |b(\tilde{X}(\infty)) - b(y)| |f'_a(y)| \right] dy \right] \\
& \leq \delta \frac{1}{\mu} \mathbb{E} \left[|b(\tilde{X}(\infty))| \right] + \delta \frac{1}{\mu} \mathbb{E} \left[|b^2(\tilde{X}(\infty)) f'_a(\tilde{X}(\infty))| \right] + \frac{1}{\mu} \mathbb{E} \left[|b^2(\tilde{X}(\infty))| \int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f'_a(y)| dy \right] \\
& \quad + 5\delta^2 \mathbb{E} \left[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)| \right] \\
& \leq \frac{10}{3} \delta + \delta \frac{1}{\mu} \mathbb{E} \left[|b^2(\tilde{X}(\infty)) f'_a(\tilde{X}(\infty))| \right] + \frac{1}{\mu} \mathbb{E} \left[|b^2(\tilde{X}(\infty))| \int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f'_a(y)| dy \right] + 5\sqrt{2}\delta^2,
\end{aligned}$$

where in the last inequality we used (5.12) and the moment bound (3.12). Now by (3.11) and (3.16),

$$\begin{aligned}
\delta \frac{1}{\mu} \mathbb{E} \left[|b^2(\tilde{X}(\infty)) f'_a(\tilde{X}(\infty))| \right] & \leq 5\delta \mathbb{E} \left[\tilde{X}^2(\infty) 1(\tilde{X}(\infty) \leq -\zeta) \right] + \delta |\zeta| \mathbb{P}(\tilde{X}(\infty) \geq -\zeta + \delta) \\
& \leq 10\delta + \delta \frac{7}{4} \leq 12\delta,
\end{aligned}$$

and similarly,

$$\frac{1}{\mu} \mathbb{E} \left[|b^2(\tilde{X}(\infty))| \int_{\tilde{X}(\infty)-\delta}^{\tilde{X}(\infty)} |f'_a(y)| dy \right] \leq 12\delta.$$

Therefore,

$$\frac{1}{\delta} \mathbb{E} \left[|b(\tilde{X}(\infty)) \epsilon_2(\tilde{X}(\infty))| \right] \leq \frac{10}{3} \delta + 24\delta + 5\sqrt{2}\delta^2 \leq 35\delta.$$

This verifies (5.11) and concludes the proof of Theorem 3. $\square$

PROOF OF THEOREM 4. The proof of this theorem is nearly identical to the proof of Theorem 3. Therefore, we only outline the key steps and differences. The goal is to obtain a version of (5.11), from which the theorem follows by applying Lemmas 10 and 11. To get a version of (5.11), we bound each of the terms in (5.8), just like we did in the proof of Theorem 3. The proof varies between the underloaded and overloaded cases.

We begin with the underloaded case ($1 \leq R \leq n$). To bound the first term in (5.8), we use moment bounds (3.18), (3.20), and (3.22), together with gradient bound (5.5). For the second and third terms in (5.8) we use the gradient bound in (5.3). For the fourth error term, we use gradient bound (5.3), and moment bounds (3.17), (3.20), and

$$\begin{aligned} \mathbb{E}\left[\left(b(\tilde{X}(\infty))\right)^2 1(\tilde{X}(\infty) \geq -\zeta)\right] &= \alpha^2 \mathbb{E}\left[\left(\tilde{X}(\infty) + \zeta\right)^2 1(\tilde{X}(\infty) \geq -\zeta)\right] + \mu^2 \zeta^2 \mathbb{P}(\tilde{X}(\infty) \geq -\zeta) \\ &\quad + 2\alpha\mu |\zeta| \mathbb{E}\left[\left(\tilde{X}(\infty) + \zeta\right) 1(\tilde{X}(\infty) \geq -\zeta)\right] \\ &\leq \alpha^2 \frac{1}{3} \left(\frac{\mu}{\alpha} \delta^2 + \frac{\mu}{\alpha} 4 + \delta^2 \right) + \mu^2 \zeta^2 \mathbb{P}(\tilde{X}(\infty) \geq -\zeta) \\ &\quad + 2\alpha\mu \left(\frac{\delta^2}{4} \frac{\alpha}{\mu} + \frac{\delta^2}{4} + 1 \right), \end{aligned}$$

where the last inequality follows from moment bounds (3.21) and (3.23).

In the overloaded case ($n \leq R$), to bound the first term in (5.8) we use moment bounds (3.25), (3.28), and (3.31) with gradient bound (5.5). To bound the second and third terms in (5.8) we use gradient bound (5.4). To bound the fourth term in (5.8), we use gradient bound (5.5), with moment bounds (3.27) and

$$\begin{aligned} \mathbb{E}\left[\left(b(\tilde{X}(\infty))\right)^2 1(\tilde{X}(\infty) \leq -\zeta)\right] &= \mu^2 \mathbb{E}\left[\left(\tilde{X}(\infty) + \zeta\right)^2 1(\tilde{X}(\infty) \leq -\zeta)\right] + \alpha^2 \zeta^2 \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) \\ &\quad + 2\alpha\mu \zeta \mathbb{E}\left[\left|\left(\tilde{X}(\infty) + \zeta\right) 1(\tilde{X}(\infty) \leq -\zeta)\right|\right] \\ &\leq \mu^2 \left(\frac{\delta^2}{4} \frac{\alpha}{\mu} + 1 \right) + \alpha^2 \left(\frac{\delta^2}{4} + \frac{\mu}{\alpha} \right) + 2\alpha\mu \left(\frac{\delta^2}{4} + 1 \right), \end{aligned}$$

where the last inequality follows from moment bounds (3.26), (3.29), and (3.30). $\square$

6. Extension: Higher Moments. In this section we discuss the use of Stein's method for the approximation of higher moments. We begin by presenting the following result for the Erlang-C model.

THEOREM 5. *Consider the Erlang-C system ($\alpha = 0$), and fix an integer $m > 0$. There exists a constant $C = C(m)$, such that for all $n \geq 1, \lambda > 0$, and $\mu > 0$ satisfying $1 \leq R < n$,*

$$(6.1) \quad \left| \mathbb{E}(\tilde{X}(\infty))^m - \mathbb{E}(Y(\infty))^m \right| \leq (1 + 1/|\zeta|^{m-1}) C(m) \delta,$$

where ζ is defined in (1.4).

The proof of this theorem follows the standard Stein framework in Section 3, but we do not provide it in this paper. The most interesting aspect of (6.1) is the appearance of $1/|\zeta|^{m-1}$ in the bound on the right hand side. This suggest that the diffusion approximation

of higher moments may not hold universally. To test this, we performed some numerical experiments illustrated in Table 3. The results suggest that the approximation error does indeed grow like $1/|\zeta|^{m-1}$, meaning that the diffusion approximation is not universal for higher moments. In [10], the authors prove error bounds for the diffusion approximation of higher moments in the Erlang-A model. They show that with α and μ fixed, the approximation error of the m th moment is bounded by $C(\alpha, \mu, m)/\sqrt{\lambda}$, where $C(\alpha, \mu, m) > 0$ is a constant that depends on μ, α , and m (they also prove a similar result when the system is in the NDS regime). In contrast to the Erlang-C model, the approximation of the Erlang-A model remains universal even for higher moments.

R	$ \zeta $	$\mathbb{E}(\tilde{X}(\infty))^2$	Error	$ \zeta \times \text{Error}$	$ \zeta ^{0.5} \times \text{Error}$	$ \zeta ^{1.5} \times \text{Error}$
499	4.48×10^{-2}	9.47×10^2	1.59	7.10×10^{-2}	0.34	1.50×10^{-2}
499.9	4.50×10^{-3}	9.94×10^4	16.50	7.38×10^{-2}	1.10	4.94×10^{-3}
499.95	2.20×10^{-3}	3.99×10^5	33.08	7.40×10^{-2}	1.56	3.50×10^{-3}
499.99	4.47×10^{-4}	9.99×10^6	165.67	7.41×10^{-2}	3.50	1.57×10^{-3}

TABLE 3

The error term above equals $|\mathbb{E}(\tilde{X}(\infty))^2 - \mathbb{E}(Y(\infty))^2|$ and grows as $R \rightarrow n$. The error term still grows when multiplied by $|\zeta|^{0.5}$, and the error term shrinks to zero when multiplied by $|\zeta|^{1.5}$. However, when multiplied by $|\zeta|$, the error term appears to converge to some limiting value, suggesting that the error does indeed grow at a rate of $1/|\zeta|$. We observed consistent behavior for higher moments of $\tilde{X}(\infty)$ as well.

Acknowledgement. This research is supported in part by NSF Grants CNS-1248117, CMMI-1335724, and CMMI-1537795. We would also like to thank the participants of the 2015 Workshop on New Directions in Stein's Method held at the Institute for Mathematical Sciences at the National University of Singapore for helpful discussions.

APPENDIX A: PROOF OF LEMMA 1

PROOF OF LEMMA 1. Let $f(x) : \mathbb{R} \rightarrow \mathbb{R}$ satisfy $|f(x)| \leq C(1+x)^2$. A sufficient condition to ensure that

$$\mathbb{E}[G_{\tilde{X}} f(\tilde{X}(\infty))] = 0$$

is given by [12, Proposition 1.1] (alternatively, see [8, Proposition 3]). Namely, we require that

$$(A.1) \quad \mathbb{E}[|G_{\tilde{X}}(\tilde{X}(\infty), \tilde{X}(\infty))f(\tilde{X}(\infty))|] < \infty,$$

where $G_{\tilde{X}}(x, x)$ is the diagonal entry of the generator matrix $G_{\tilde{X}}$ corresponding to state x .

We begin with the Erlang-C model, where the transition rates of the CTMC are bounded by $\lambda + n\mu$. Since $|f(x)| \leq C(1+x)^2$, it suffices to show that $\mathbb{E}(\tilde{X}(\infty))^2 < \infty$, or that $\mathbb{E}(X(\infty))^2 < \infty$, where $X(\infty)$ has the stationary distribution of the CTMC X . Consider the function $V(k) = k^3$, where $k \in \mathbb{Z}_+$. Let G_X be the generator of X , which is a simple

birth death process with constant birth rate λ and departure rate $\mu(k \wedge n)$ in state $k \in \mathbb{Z}_+$. Then for $k \geq n$,

$$\begin{aligned}
 G_X V(k) &= \lambda((k+1)^3 - k^3) + n\mu((k-1)^3 - k^3) \\
 &= \lambda(3k^2 + 3k + 1) + n\mu(-3k^2 + 3k - 1) \\
 (A.2) \quad &= -3k^2(n\mu - \lambda) + 3k(\lambda + n\mu) + (\lambda - n\mu).
 \end{aligned}$$

It is not hard to see that there exists some $k_0 \in \mathbb{Z}_+$, and a constant $c > 0$ (that depends on λ, n , and μ), such that for all $k \geq k_0$,

$$(A.3) \quad -3k^2(n\mu - \lambda) + 3k(\lambda + n\mu) \leq -ck^2.$$

We combine (A.2)–(A.3) to conclude that there exists some constant $d > 0$ (that depends on λ, n , and μ) satisfying

$$G_X V(x) \leq -cx^2 + d1(k < (k_0 \vee n)),$$

and invoking [13, Theorem 4.3], we see that $\mathbb{E}(X(\infty))^2 < \infty$.

The case of the Erlang-A model is not very different. When $\alpha > 0$, the transition rates of the CTMC depend linearly on its state. Hence, to satisfy (A.1) we need to show that $\mathbb{E}(X(\infty))^3 < \infty$. This is readily proven by repeating the procedure above with the Lyapunov function $V(k) = k^4$, and we omit the details. $\square$

APPENDIX B: MOMENT BOUNDS

B.1. Proof of Lemma 2.

PROOF OF LEMMA 2. We first prove (3.11), (3.12), and (3.14). Recalling the generator $G_{\tilde{X}}$ defined in (3.6), we apply it to the function $V(x) = x^2$ to see that for $k \in \mathbb{Z}_+$ and $x = x_k = \delta(k - x(\infty))$,

$$\begin{aligned}
 G_{\tilde{X}} V(x) &= \lambda(2x\delta + \delta^2) + \mu(k \wedge n)(-2x\delta + \delta^2) \\
 &= 2x\delta(\lambda - n\mu + \mu(k - n)^-) + \mu + \delta^2\mu(k \wedge n) \\
 &= 2x\mu(\zeta + (x + \zeta)^-) + \mu + \delta^2\mu(n - \frac{\lambda}{\mu} + \frac{\lambda}{\mu} - (k - n)^-) \\
 &= 2x\mu(\zeta + (x + \zeta)^-) + \mu - \delta\mu\zeta + \mu - \delta\mu(x + \zeta)^- \\
 &= 1(x \leq -\zeta)\mu(-2x^2 + \delta x) + 1(x > -\zeta)\mu(2x\zeta - \delta\zeta) + 2\mu \\
 (B.1) \quad &\leq 1(x \leq -\zeta)\mu(-\frac{3}{2}x^2 + \frac{\delta^2}{2}) + 1(x > -\zeta)\mu(2x\zeta - \delta\zeta) + 2\mu.
 \end{aligned}$$

Instead of splitting the last two lines into the cases $x \leq -\zeta$ and $x > -\zeta$, we could have also considered $x < -\zeta$ and $x \geq -\zeta$ instead, and would have obtained

$$\begin{aligned} G_{\tilde{X}}V(x) &= 1(x < -\zeta)\mu(-2x^2 + \delta x) + 1(x \geq -\zeta)\mu(2x\zeta - \delta\zeta) + 2\mu \\ (B.2) \quad &\leq 1(x < -\zeta)\mu\left(-\frac{3}{2}x^2 + \frac{\delta^2}{2}\right) + 1(x \geq -\zeta)\mu(2x\zeta - \delta\zeta) + 2\mu. \end{aligned}$$

We take expected values on both sides of (B.1) with respect to $\tilde{X}(\infty)$, and apply Lemma 1 to see that

$$\begin{aligned} 0 &\leq -\frac{3}{2}\mu\mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)] \\ (B.3) \quad &+ \mu|\zeta|\mathbb{E}[(-2\tilde{X}(\infty) + \delta)1(\tilde{X}(\infty) > -\zeta)] + 2\mu + \frac{\mu\delta^2}{2}. \end{aligned}$$

This implies that when $|\zeta| > \delta/2$,

$$0 \leq -\frac{3}{2}\mu\mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)] + 2\mu + \frac{\mu\delta^2}{2},$$

and when $|\zeta| \leq \delta/2$,

$$0 \leq -\frac{3}{2}\mu\mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)] + 2\mu + \mu\delta^2.$$

Therefore,

$$\mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)] \leq \frac{4}{3} + \frac{2\delta^2}{3},$$

which proves (3.11). Jensen's inequality immediately gives us

$$\mathbb{E}[|\tilde{X}(\infty)1(\tilde{X}(\infty) \leq -\zeta)|] \leq \sqrt{\mathbb{E}[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \leq -\zeta)]},$$

which proves (3.12). Furthermore, (B.3) also gives us

$$\mathbb{E}[|\tilde{X}(\infty)1(\tilde{X}(\infty) > -\zeta)|] \leq \frac{1}{|\zeta|} + \frac{\delta^2}{4|\zeta|} + \frac{\delta}{2},$$

which is not quite (3.14) because the inequality above has $1(\tilde{X}(\infty) > -\zeta)$ as opposed to $1(\tilde{X}(\infty) \geq -\zeta)$ as in (3.14). However, we can use (B.2) to get the stronger bound

$$\mathbb{E}[|\tilde{X}(\infty)1(\tilde{X}(\infty) \geq -\zeta)|] \leq \frac{1}{|\zeta|} + \frac{\delta^2}{4|\zeta|} + \frac{\delta}{2},$$

which proves (3.14).

We now prove (3.13), or

$$(B.4) \quad \mathbb{E} \left[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)| \right] \leq 2|\zeta|.$$

We use the triangle inequality to see that

$$\mathbb{E} \left[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \leq -\zeta)| \right] \leq |\zeta| + \mathbb{E} \left[|\tilde{X}(\infty) + \zeta| 1(\tilde{X}(\infty) \leq -\zeta) \right].$$

The second term on the right hand side is just expected number of idle servers, scaled by δ . We now show that this expected value equals $|\zeta|$. Applying the generator $G_{\tilde{X}}$ to the test function $f(x) = x$, one sees that for all $k \in \mathbb{Z}_+$ and $x = x_k = \delta(k - x(\infty))$,

$$G_{\tilde{X}}f(x) = \delta\lambda - \delta\mu(k \wedge n) = \mu[\zeta + (x + \zeta)^-].$$

Taking expected values with respect to $\tilde{X}(\infty)$ on both sides, and applying Lemma 1, we arrive at

$$(B.5) \quad \mathbb{E} \left[|(\tilde{X}(\infty) + \zeta) 1(\tilde{X}(\infty) \leq -\zeta)| \right] = |\zeta|,$$

which proves (3.13).

We move on to prove (3.15), or

$$(B.6) \quad \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) \leq (2 + \delta) |\zeta|.$$

Let I be the unscaled expected number of idle servers. Then by (B.5),

$$I = \mathbb{E}(X(\infty) - n)^- = \frac{1}{\delta} \mathbb{E} \left[|(\tilde{X}(\infty) + \zeta) 1(\tilde{X}(\infty) \leq -\zeta)| \right] = \frac{1}{\delta} |\zeta|.$$

Now let $\{\nu_k\}_{k=0}^\infty$ be the stationary distribution of X (the unscaled CTMC). We want to prove an upper bound on the probability

$$\mathbb{P}(\tilde{X}(\infty) \leq -\zeta) = \sum_{k=0}^n \nu_k \leq \sum_{k=0}^{\lfloor n-\sqrt{R} \rfloor} \nu_k + \sum_{k=\lceil n-\sqrt{R} \rceil}^n \nu_k.$$

Observe that

$$I = \sum_{k=0}^n (n-k) \nu_k \geq \sqrt{R} \sum_{k=0}^{\lfloor n-\sqrt{R} \rfloor} \nu_k.$$

Now let k^* be the first index that maximizes $\{\nu_k\}_{k=0}^\infty$, i.e.

$$k^* = \inf\{k \geq 0 : \nu_k \geq \nu_j, \text{ for all } j \neq k\}.$$

Then

$$\begin{aligned} \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) &= \sum_{k=0}^{\lfloor n-\sqrt{R} \rfloor} \nu_k + \sum_{k=\lceil n-\sqrt{R} \rceil}^n \nu_k \leq \frac{I}{\sqrt{R}} + (\sqrt{R}+1)\nu_{k^*} \\ (B.7) \qquad \qquad \qquad &= |\zeta| + (\sqrt{R}+1)\nu_{k^*}. \end{aligned}$$

Applying $G_{\tilde{X}}$ to the test function $f(x) = (k \wedge k^*)$, we see that for all $k \in \mathbb{Z}_+$ and $x = x_k = \delta(k - x(\infty))$,

$$G_{\tilde{X}}f(x) = \delta\lambda 1(k < k^*) - \delta\mu(k \wedge n)1(k \leq k^*).$$

Taking expected values with respect to $X(\infty)$ on both sides and applying Lemma 1, we see that

$$\mathbb{P}(X(\infty) \leq k^*) = \frac{\mu}{n\mu - \lambda} \mathbb{E}[(X(\infty) - n)^{-1} 1(X(\infty) \leq k^*)] - \nu_{k^*} \frac{\lambda}{n\mu - \lambda} \geq 0.$$

Using the inequality above, together with the fact that $k^* \leq n$, we see that

$$\nu_{k^*} \leq \frac{\mu}{\lambda} \mathbb{E}[(X(\infty) - n)^{-1} 1(X(\infty) \leq k^*)] \leq \frac{\mu}{\lambda} \mathbb{E}[(X(\infty) - n)^{-1} 1(X(\infty) \leq n)] = \frac{I}{R} = \frac{|\zeta|}{\sqrt{R}}.$$

The fact that $k^* \leq n$ is a consequence of $\lambda < n\mu$, and can be verified through the flow balance equations of the CTMC X . We combine the bound above with (B.7) to arrive at (3.15), which concludes the proof of this lemma. $\square$

B.2. Proof Outline for Lemma 3: The Underloaded System. The proof of the underloaded case of Lemma 3 is very similar to that of Lemma 2. Therefore, we only outline some key intermediate steps needed to obtain the results. We remind the reader that when $R \leq n$, then $\zeta \leq 0$. We first show how to establish (3.17), which is proved in a similar fashion to (3.11) of Lemma 2 – by applying the generator $G_{\tilde{X}}$ to the Lyapunov function $V(x) = x^2$. The following are some useful intermediate steps for any reader wishing to produce a complete proof. The first step to prove (3.17) is to get an analogue of (B.1). Namely, when $x \leq -\zeta$,

$$G_{\tilde{X}}V(x) = -2\mu x^2 + \mu\delta x + 2\mu \leq -\frac{3}{2}\mu x^2 + \mu\delta^2/2 + 2\mu,$$

and when $x \geq -\zeta$,

$$\begin{aligned} G_{\tilde{X}}V(x) &= -2\alpha(x + \zeta)^2 + \alpha\delta(x + \zeta) - 2\mu|\zeta|(x + \zeta) - 2|\zeta|\alpha(x + \zeta) + \mu|\zeta|(\delta - 2|\zeta|) + 2\mu \\ (B.8) \quad &\leq -\frac{3}{2}\alpha(x + \zeta)^2 - 2\mu|\zeta|(x + \zeta) + \delta^2\alpha/2 + \delta^2\mu/8 + 2\mu. \end{aligned}$$

From here, we use Lemma 1 to get a statement similar to (B.3), from which we can infer (3.17) and by applying Jensen's inequality to (3.17), we get (3.18). Observe that this procedure yields (3.21), (3.22), and (3.23) as well. We now describe how to prove (3.20), which requires only a slight modification of (B.8). Namely, for $x \geq -\zeta$,

$$G_{\tilde{X}}V(x) = 2x(-\alpha(x + \zeta) + \mu\zeta) - \delta(-\alpha(x + \zeta) + \mu\zeta) + 2\mu.$$

From this, we can deduce that since $x \geq -\zeta$,

$$G_{\tilde{X}}V(x) \leq -2(\mu \wedge \alpha)x^2 - \delta(-\alpha(x + \zeta) + \mu\zeta) + 2\mu,$$

and also

$$G_{\tilde{X}}V(x) \leq -2\mu|\zeta|x - \delta(-\alpha(x + \zeta) + \mu\zeta) + 2\mu.$$

Then Lemma 1 can be applied as before to see that both

$$(B.9) \quad 2\mu|\zeta|\mathbb{E}\left[|\tilde{X}(\infty)1(\tilde{X}(\infty) \geq -\zeta)|\right] \quad \text{and} \quad 2(\mu \wedge \alpha)\mathbb{E}\left[(\tilde{X}(\infty))^2 1(\tilde{X}(\infty) \geq -\zeta)\right]$$

are bounded by

$$2\mu + \mu\delta^2/2 - \delta\mathbb{E}\left[(-\alpha(\tilde{X}(\infty) + \zeta) + \mu\zeta)1(\tilde{X}(\infty) \geq -\zeta)\right].$$

Applying the generator $G_{\tilde{X}}$ to the test function $f(x) = x$ and taking expected values with respect to $\tilde{X}(\infty)$, we get $\mathbb{E}b(\tilde{X}(\infty)) = 0$, or

$$(B.10) \quad \mathbb{E}\left[(-\alpha(\tilde{X}(\infty) + \zeta) + \mu\zeta)1(\tilde{X}(\infty) \geq -\zeta)\right] = \mu\mathbb{E}\left[\tilde{X}(\infty)1(\tilde{X}(\infty) < -\zeta)\right].$$

When combined with (3.18), this implies that

$$\begin{aligned} & 2\mu + \mu\delta^2/2 - \delta\mathbb{E}\left[(-\alpha(\tilde{X}(\infty) + \zeta) + \mu\zeta)1(\tilde{X}(\infty) \geq -\zeta)\right] \\ & \leq 2\mu + \mu\delta^2/2 + \mu\delta\sqrt{\frac{1}{3}\left(\frac{\alpha}{\mu}\delta^2 + \delta^2 + 4\right)}, \end{aligned}$$

which proves (3.20), because the quantity above is an upper bound for (B.9). To prove (3.19), we manipulate (B.10) to get

$$\mathbb{E}\left[|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) \leq -\zeta)|\right] = |\zeta| + \frac{\alpha}{\mu}\mathbb{E}\left[|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) > -\zeta)|\right],$$

to which we can apply the triangle inequality and (3.22) to conclude (3.19). Lastly, the proof of (3.24) is nearly identical to the proof of (3.15) in Lemma 2. The key step is to obtain an analogue of (B.7).

B.3. Proof Outline for Lemma 3: The Overloaded System. The proof of the overloaded case of Lemma 3 is also similar to that of Lemma 2. Therefore, we only outline some key intermediate steps needed to obtain the results; the bounds in this lemma are not proved in the order in which they are stated. We remind the reader that when $R \geq n$, then $\zeta \geq 0$. We start by proving (3.27). Although the left hand side of (3.27) is slightly different from (3.11) of Lemma 2, it is proved using the same approach – by applying the generator $G_{\tilde{X}}$ to the Lyapunov function $V(x) = x^2$. The following are some useful intermediate steps for any reader wishing to produce a complete proof. The first step to prove (3.27) is to get analogue of (B.1). Namely, when $x \leq -\zeta$,

$$\begin{aligned} G_{\tilde{X}}V(x) &= -2\mu(x + \zeta)^2 + \mu\delta(x + \zeta) + 2(\mu + \alpha)|\zeta|(x + \zeta) - 2\alpha\zeta^2 - \alpha\delta\zeta + 2\mu \\ (B.11) \quad &\leq -2\mu(x + \zeta)^2 + 2(\mu + \alpha)|\zeta|(x + \zeta) + 2\mu, \end{aligned}$$

and when $x \geq -\zeta$,

$$G_{\tilde{X}}V(x) = -2\alpha x^2 + \alpha\delta x + 2\mu \leq -\frac{3}{2}\alpha x^2 + \alpha\delta^2/2 + 2\mu.$$

From here, we use Lemma 1 to get a statement similar to (B.3), which implies (3.27). Applying Jensen's inequality to (3.27) yields (3.28). The procedure used to get (3.27) also yields (3.29), (3.30), and (3.31).

We now describe how to prove (3.25) and (3.26), which requires only a slight modification of (B.11). Namely, we use the fact that for $x \leq -\zeta$,

$$G_{\tilde{X}}V(x) = 2x(-\mu(x + \zeta) + \alpha\zeta) - \delta(-\mu(x + \zeta) + \alpha\zeta) + 2\mu.$$

From this, one can deduce that since $x \leq -\zeta$,

$$G_{\tilde{X}}V(x) \leq -2(\mu \wedge \alpha)x^2 + 2\mu,$$

and also

$$G_{\tilde{X}}V(x) \leq -2\alpha|\zeta||x| + 2\mu.$$

Then Lemma 1 and Jensen's inequality can be applied as before to get both (3.25) and (3.26).

We now prove (3.32). Observe that

$$\begin{aligned} \mathbb{E}\left[\left|\tilde{X}(\infty)1(\tilde{X}(\infty) \geq -\zeta)\right|\right] &= \mathbb{E}\left[\left|(\tilde{X}(\infty) + \zeta - \zeta)1(\tilde{X}(\infty) \geq -\zeta)\right|\right] \\ &\geq \mathbb{E}\left[\left|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) > -\zeta) - \zeta 1(\tilde{X}(\infty) > -\zeta)\right|\right] \\ &\geq \mathbb{E}\left[\left|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) > -\zeta)\right|\right] - \zeta \\ &= \frac{\mu}{\alpha}\mathbb{E}\left[\left|(\tilde{X}(\infty) + \zeta)1(\tilde{X}(\infty) \leq -\zeta)\right|\right], \end{aligned}$$

where the last equality comes from applying the generator $G_{\tilde{X}}$ to the function $f(x) = x$ and taking expected values with respect to $\tilde{X}(\infty)$ to see that $\mathbb{E}b(\tilde{X}(\infty)) = 0$, or

$$(B.12) \quad \mathbb{E} \left[(-\mu(\tilde{X}(\infty) + \zeta) + \alpha\zeta) 1(\tilde{X}(\infty) \leq -\zeta) \right] = \alpha \mathbb{E} \left[\tilde{X}(\infty) 1(\tilde{X}(\infty) > -\zeta) \right].$$

Therefore,

$$\mathbb{E} \left[|(\tilde{X}(\infty) + \zeta) 1(\tilde{X}(\infty) \leq -\zeta)| \right] \leq \frac{\alpha}{\mu} \mathbb{E} \left[|\tilde{X}(\infty) 1(\tilde{X}(\infty) \geq -\zeta)| \right],$$

and we can invoke (3.28) to conclude (3.32).

We now prove (3.33), which requires additional arguments that we have not used in the proof of Lemma 2. We assume for now that

$$(B.13) \quad \lambda \leq n\mu + \frac{1}{2}\sqrt{n}\mu.$$

Fix $\gamma \in (0, 1/2)$, and define

$$(B.14) \quad J_1 = \sum_{k=0}^{\lfloor n-\gamma\sqrt{R} \rfloor} \nu_k, \quad J_2 = \sum_{k=\lceil n-\gamma\sqrt{R} \rceil}^n \nu_k,$$

where $\{\nu_k\}_{k=0}^{\infty}$ is the stationary distribution of X . We note that by (B.13),

$$n/\sqrt{R} \geq \sqrt{R} - \frac{1}{2}\sqrt{n/R} \geq \sqrt{R} - 1/2 \geq 1/2,$$

which implies that $n - \gamma\sqrt{R} > 0$. Then $\mathbb{P}(\tilde{X}(\infty) \leq -\zeta) = \mathbb{P}(X(\infty) \leq n) \leq J_1 + J_2$. To bound J_1 we observe that

$$\mathbb{E} \left[\left| \tilde{X}(\infty) + \zeta \right| 1_{\{\tilde{X}(\infty) \leq -\zeta\}} \right] = \frac{1}{\sqrt{R}} \sum_{k=0}^n (n-k) \nu_k \geq \gamma \sum_{k=0}^{\lfloor n-\gamma\sqrt{R} \rfloor} \nu_k = \gamma J_1.$$

Combining (3.29)–(3.32), we conclude that

$$(B.15) \quad \begin{aligned} J_1 &\leq \frac{1}{\gamma} \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \vee 1 \wedge \frac{\alpha}{\mu} \sqrt{\frac{\mu}{\alpha}} \vee 1 \right) \\ &\leq \frac{1}{\gamma} \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right). \end{aligned}$$

Now to bound J_2 , we apply $G_{\tilde{X}}$ to the test function $f(x) = k \wedge n$, where $x = \delta(k - x(\infty))$, and take the expectation with respect to $\tilde{X}(\infty)$ to see that

$$0 = -\lambda\nu_n + (\lambda - n\mu)\mathbb{P}(X(\infty) \leq n) + \mu \mathbb{E} \left[(X(\infty) - n)^- 1_{\{X(\infty) \leq n\}} \right].$$

Noticing that

$$\mathbb{E}[(X(\infty) - n)^- 1_{\{X(\infty) \leq n\}}] = \frac{1}{\delta} \mathbb{E} \left[\left| \tilde{X}(\infty) + \zeta \right| 1_{\{\tilde{X}(\infty) \leq -\zeta\}} \right],$$

we arrive at

$$(B.16) \quad \nu_n \leq \delta \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) + \frac{\lambda - n\mu}{\lambda} \mathbb{P}(X(\infty) \leq n).$$

The flow balance equations

$$\lambda \nu_{k-1} = k\mu \nu_k, \quad k = 1, 2, \dots, n$$

imply that $\nu_0 < \nu_1 < \dots < \nu_{n-2} < \nu_{n-1} \leq \nu_n$, and therefore

$$(B.17) \quad \begin{aligned} J_2 &\leq (\gamma\sqrt{R} + 1)\nu_n \leq (\gamma\sqrt{R} + 1) \left[\delta \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) + \frac{\lambda - n\mu}{\lambda} \mathbb{P}(X(\infty) \leq n) \right] \\ &= (\gamma + \delta) \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) \\ &\quad + (\gamma\sqrt{R} + 1) \frac{\lambda - n\mu}{\lambda} J_1 + (\gamma\sqrt{R} + 1) \frac{\lambda - n\mu}{\lambda} J_2 \end{aligned}$$

We use (B.13), the fact that $\gamma \in (0, 1/2)$, and that $R \geq n \geq 1$ to see that

$$(\gamma\sqrt{R} + 1) \frac{\lambda - n\mu}{\lambda} \leq (\gamma\sqrt{R} + 1) \frac{\sqrt{n}}{2R} \leq \frac{1}{2}(\gamma + 1/\sqrt{R}) = \frac{1}{2}(\gamma + 1) < 3/4.$$

Then by rearranging terms in (B.17) and applying (B.15) we conclude that

$$\begin{aligned} \frac{1}{4} J_2 &\leq (\gamma + \delta) \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) + \frac{3}{4} \frac{1}{\gamma} \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) \\ &= \left(\gamma + \delta + \frac{3}{4} \frac{1}{\gamma} \right) \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right). \end{aligned}$$

Hence, we have just shown that under assumption (B.13),

$$\begin{aligned} \mathbb{P}(\tilde{X}(\infty) \leq -\zeta) &\leq J_1 + J_2 \leq \frac{1}{\gamma} \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) \\ &\quad + 4 \left(\gamma + \delta + \frac{3}{4} \frac{1}{\gamma} \right) \frac{2}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right) \\ &\leq (3 + \delta) \frac{8}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta} \wedge \sqrt{\frac{\alpha}{\mu}} \right), \end{aligned}$$

where to get the last inequality we fixed $\gamma \in (0, 1/2)$ that solves $\gamma + 1/\gamma = 3$.

We now wish to establish the same result without assumption (B.13), i.e. when $\lambda > n\mu + \frac{1}{2}\sqrt{n\mu}$. For this, we rely on the following comparison result. Fix n, μ and α and let $X^{(\lambda)}(\infty)$ be the steady-state customer count in an Erlang-A system with arrival rate λ , service rate μ , number of servers n , and abandonment rate α . Then for any $0 < \lambda_1 < \lambda_2$,

$$(B.18) \quad \mathbb{P}(X^{(\lambda_2)}(\infty) \leq n) \leq \mathbb{P}(X^{(\lambda_1)}(\infty) \leq n).$$

This says that with all other parameters being held fixed, an Erlang-A system with a higher arrival rate is less likely to have idle servers. Intuitively, this statement is obvious. To prove it, one can use the flow balance equations to see that

$$\begin{aligned} \mathbb{P}(X^{(\lambda)}(\infty) \leq n) &= \frac{\sum_{j=0}^n \frac{(\lambda/\mu)^j}{j!}}{\sum_{j=0}^n \frac{(\lambda/\mu)^j}{j!} + \sum_{j=n+1}^{\infty} \frac{(\lambda/\mu)^n}{n!} \prod_{k=n+1}^j \frac{\lambda}{n\mu + (k-n)\alpha}} \\ &= \frac{1}{1 + \frac{\sum_{j=n+1}^{\infty} \frac{(\lambda/\mu)^n}{n!} \prod_{k=n+1}^j \frac{\lambda}{n\mu + (k-n)\alpha}}{\sum_{j=0}^n \frac{(\lambda/\mu)^j}{j!}}}. \end{aligned}$$

Now

$$\frac{\sum_{j=n+1}^{\infty} \frac{(\lambda/\mu)^n}{n!} \prod_{k=n+1}^j \frac{\lambda}{n\mu + (k-n)\alpha}}{\sum_{j=0}^n \frac{(\lambda/\mu)^j}{j!}} = \frac{\sum_{j=n+1}^{\infty} \frac{(1/\mu)^n}{n!} \prod_{k=n+1}^j \frac{\lambda}{n\mu + (k-n)\alpha}}{\sum_{j=0}^n \frac{1}{\lambda^{n-j}} \frac{(1/\mu)^j}{j!}},$$

is an increasing function of λ , which implies (B.18). Therefore, for $\lambda > n\mu + \frac{1}{2}\sqrt{n\mu}$,

$$\mathbb{P}(X^{(\lambda)}(\infty) \leq n) \leq \mathbb{P}(X^{(n\mu + \frac{1}{2}\sqrt{n\mu})}(\infty) \leq n) \leq (3 + \delta) \frac{8}{\sqrt{3}} \left(\frac{\delta^2}{4} + 1 \right) \left(\frac{1}{\zeta^{(n\mu + \frac{1}{2}\sqrt{n\mu})}} \wedge \sqrt{\frac{\alpha}{\mu}} \right)$$

where $\zeta^{(n\mu + \frac{1}{2}\sqrt{n\mu})}$ is the ζ corresponding to $X^{(n\mu + \frac{1}{2}\sqrt{n\mu})}(\infty)$, and satisfies

$$\frac{1}{\zeta^{(n\mu + \frac{1}{2}\sqrt{n\mu})}} = \frac{2\alpha}{\mu} \sqrt{\frac{n + \sqrt{n}/2}{n}} \leq \frac{2\alpha}{\mu} \sqrt{\frac{3}{2}}.$$

This concludes the proof of (3.33).

APPENDIX C: GRADIENT BOUNDS

In this section we prove the gradient bounds in Lemmas 4–7. The proofs of these lemmas follows the arguments of the proof of [6, Lemma 13.1]. For completeness, we repeat many of those arguments in this section. We recall the Poisson equation (3.3), or

$$G_Y f_h(x) = \mu f_h''(x) + b(x) f_h'(x) = \mathbb{E}h(Y(\infty)) - h(x), \quad x \in \mathbb{R}, \quad h(x) \in \mathcal{H}.$$

Furthermore, recall that $\nu(x)$ is the density of $Y(\infty)$, and satisfies

$$\nu(x) = \kappa \exp\left(\frac{1}{\mu} \int_0^x b(y) dy\right),$$

where κ is a normalizing constant. Now recall that the family of solutions to the Poisson equation is given by (3.4), and is parametrized by constants $a_1, a_2 \in \mathbb{R}$. We fix a solution $f_h(x)$ with $a_2 = 0$, and see that for this solution

$$(C.1) \quad f'_h(x) = \frac{1}{\nu(x)} \int_{-\infty}^x \frac{1}{\mu} (\mathbb{E}h(Y(\infty)) - h(y)) \nu(y) dy, \quad x \in \mathbb{R}.$$

Observe that since $\nu(x)$ is the density of $Y(\infty)$,

$$\int_{-\infty}^{\infty} \frac{1}{\mu} (\mathbb{E}h(Y(\infty)) - h(y)) \nu(y) dy = 0,$$

which implies that

$$(C.2) \quad f'_h(x) = - \frac{1}{\nu(x)} \int_x^{\infty} \frac{1}{\mu} (\mathbb{E}h(Y(\infty)) - h(y)) \nu(y) dy.$$

We will see that to establish gradient bounds, we will have to make use of both expressions for $f'_h(x)$ in (C.1) and (C.2), depending on the value of x . It is here that the relationship between the diffusion process Y and the random variable $Y(\infty)$ surfaces. If $\mathbb{E}h(Y(\infty))$ were to be replaced by any other constant, then (C.2) would not hold. The reason $\mathbb{E}h(Y(\infty))$ is the “correct” constant is because $Y(\infty)$ has the stationary distribution of the diffusion process Y .

We now proceed to prove the gradient bounds. First in case when $\mathcal{H} = \text{Lip}(1)$ (the Wasserstein setting), and then when $\mathcal{H} = \mathcal{H}_K$ (the Kolmogorov setting).

C.1. Wasserstein Gradient Bounds: Proof of Lemmas 4 and 5 . Fix $h(x) \in \text{Lip}(1)$; without loss of generality we assume that $h(0) = 0$. We now derive some equations that will be useful to prove both Lemmas 4 and 5. From the form of $f'_h(x)$ in (C.1) and (C.2), we have

$$\begin{aligned} f'_h(x) &\leq \frac{1}{\mu\nu(x)} \int_{-\infty}^x |y| \nu(y) dy + \frac{\mathbb{E}|Y(\infty)|}{\mu\nu(x)} \int_{-\infty}^x \nu(y) dy, \\ f'_h(x) &\leq \frac{1}{\mu\nu(x)} \int_x^{\infty} |y| \nu(y) dy + \frac{\mathbb{E}|Y(\infty)|}{\mu\nu(x)} \int_x^{\infty} \nu(y) dy. \end{aligned}$$

Now by differentiating the Poisson equation (3.3), we see that for those $x \in \mathbb{R}$ where $h'(x)$ and $b'(x)$ are defined,

$$f_h'''(x) = \frac{1}{\mu} [-h'(x) - b'(x)f'_h(x) - b(x)f_h''(x)].$$

By observing that $b(x) = \mu\nu'(x)/\nu(x)$, we can rearrange the terms above and multiply both sides by $\nu(x)$ to get

$$(C.3) \quad \frac{\nu(x)}{\mu}(-h'(x) - f'_h(x)b'(x)) = \nu(x)f_h'''(x) + \nu'(x)f_h''(x) = (f_h''(x)\nu(x))'.$$

Suppose we know that

$$(C.4) \quad \lim_{x \rightarrow \pm\infty} \nu(x)f_h''(x) = 0,$$

a fact that will be verified separately for each of Lemmas 4 and 5, right after bounding $f_h'(x)$ in the proofs of those lemmas. Since $\lim_{x \rightarrow -\infty} \nu(x)f_h''(x) = 0$, we integrate (C.3) to get

$$f_h''(x) = \frac{1}{\nu(x)} \int_{-\infty}^x \frac{1}{\mu}(-h'(y) - f'_h(y)b'(y))\nu(y)dy,$$

and since $\lim_{x \rightarrow \infty} \nu(x)f_h''(x) = 0$, it is also true that

$$f_h''(x) = -\frac{1}{\nu(x)} \int_x^{\infty} \frac{1}{\mu}(-h'(y) - f'_h(y)b'(y))\nu(y)dy.$$

To summarize, we have just shown that provided (C.4) holds,

$$(C.5) \quad |f'_h(x)| \leq \frac{1}{\mu\nu(x)} \int_{-\infty}^x |y| \nu(y)dy + \frac{\mathbb{E}|Y(\infty)|}{\mu\nu(x)} \int_{-\infty}^x \nu(y)dy,$$

$$(C.6) \quad |f'_h(x)| \leq \frac{1}{\mu\nu(x)} \int_x^{\infty} |y| \nu(y)dy + \frac{\mathbb{E}|Y(\infty)|}{\mu\nu(x)} \int_x^{\infty} \nu(y)dy,$$

$$(C.7) \quad f_h''(x) = \frac{1}{\nu(x)} \int_{-\infty}^x \frac{1}{\mu}(-h'(y) - f'_h(y)b'(y))\nu(y)dy$$

$$(C.8) \quad = -\frac{1}{\nu(x)} \int_x^{\infty} \frac{1}{\mu}(-h'(y) - f'_h(y)b'(y))\nu(y)dy,$$

$$(C.9) \quad f_h'''(x) = \frac{1}{\mu}[-h'(x) - f_h''(x)b(x) - f'_h(x)b'(x)],$$

where $f_h'''(x)$ is defined for all $x \in \mathbb{R}$ such that $h'(x)$ and $b'(x)$ exist. The multiple bounds for $f_h'(x)$ are convenient, because we can choose which one to use based on the value of x . For example, suppose $\zeta \leq 0$. Then when $x \leq 0$, we will use (C.5), when $x \geq -\zeta$ we will use (C.6), and when $x \in [0, -\zeta]$ we will use both (C.5) and (C.6) and choose the better bound. The same applies to $f_h''(x)$.

C.1.1. *Proof of Lemma 4.* The following lemma presents several bounds that will be used to prove Lemma 4. We prove it at the end of this section.

LEMMA 12. *Consider the Erlang-C model ($\alpha = 0$), and let $\nu(x)$ be the density of $Y(\infty)$. Then*

$$(C.10) \quad \frac{1}{\nu(x)} \int_{-\infty}^x \nu(y) dy \leq \begin{cases} \sqrt{\frac{\pi}{2}}, & x \leq 0, \\ \sqrt{2\pi} e^{\frac{1}{2}\zeta^2}, & x \in [0, -\zeta], \end{cases}$$

$$(C.11) \quad \frac{1}{\nu(x)} \int_x^{\infty} \nu(y) dy \leq \begin{cases} \sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|}, & x \in [0, -\zeta], \\ \frac{1}{|\zeta|}, & x \geq -\zeta, \end{cases}$$

$$(C.12) \quad \frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy \leq \begin{cases} 1, & x \leq 0, \\ 2e^{\frac{1}{2}\zeta^2} - 1, & x \in [0, -\zeta], \end{cases}$$

$$(C.13) \quad \frac{1}{\nu(x)} \int_x^{\infty} |y| \nu(y) dy \leq \begin{cases} 2 + \frac{1}{\zeta^2}, & x \in [0, -\zeta], \\ \frac{x}{|\zeta|} + \frac{1}{\zeta^2}, & x \geq -\zeta, \end{cases}$$

$$(C.14) \quad \frac{|b(x)|}{\mu\nu(x)} \int_{-\infty}^x \nu(y) dy \leq 1, \quad x \leq 0,$$

$$(C.15) \quad \frac{|b(x)|}{\mu\nu(x)} \int_x^{\infty} \nu(y) dy \leq 2, \quad x \geq 0,$$

$$(C.16) \quad \mathbb{E} |Y(\infty)| \leq \frac{1}{|\zeta|} + 1.$$

PROOF OF LEMMA 4. We begin by bounding $f'_h(x)$ by applying Lemma 12 to (C.5) and (C.6). For $x \leq -\zeta$, we apply (C.10), (C.12), and (C.16) to (C.5), and for $x \geq 0$, we apply (C.11), (C.13), and (C.16) to (C.6) to see that

$$(C.17) \quad \begin{aligned} \mu |f'_h(x)| &\leq 1 + \sqrt{\frac{\pi}{2}} \left(1 + \frac{1}{|\zeta|}\right) \leq 2.3 + 1.3/|\zeta|, \quad x \leq 0, \\ \mu |f'_h(x)| &\leq \min \left\{ 2e^{\frac{1}{2}\zeta^2} - 1 + \sqrt{2\pi} e^{\frac{1}{2}\zeta^2} \left(1 + \frac{1}{|\zeta|}\right), \right. \\ &\quad \left. 2 + \frac{1}{\zeta^2} + \left(\sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|}\right) \left(1 + \frac{1}{|\zeta|}\right) \right\}, \quad x \in [0, -\zeta], \\ \mu |f'_h(x)| &\leq \frac{x}{|\zeta|} + \frac{1}{\zeta^2} + \frac{1}{|\zeta|} \left(1 + \frac{1}{|\zeta|}\right) \leq \frac{1}{|\zeta|} (x + 1 + 2/|\zeta|), \quad x \geq -\zeta. \end{aligned}$$

For $x \in [0, -\zeta]$, observe that when $|\zeta| \leq 1$, then

$$2e^{\frac{1}{2}\zeta^2} - 1 + \sqrt{2\pi} e^{\frac{1}{2}\zeta^2} \left(1 + \frac{1}{|\zeta|}\right) \leq 3.3 - 1 + 4.2 \left(1 + \frac{1}{|\zeta|}\right) = 6.5 + 4.2/|\zeta|,$$

and when $|\zeta| \geq 1$, then

$$2 + \frac{1}{\zeta^2} + \left(\sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|} \right) \left(1 + \frac{1}{|\zeta|} \right) \leq 3 + (1.3 + 1) \left(1 + \frac{1}{|\zeta|} \right) = 5.3 + 2.3/|\zeta|.$$

Therefore,

$$(C.18) \quad |f'_h(x)| \leq \begin{cases} \frac{1}{\mu}(6.5 + 4.2/|\zeta|), & x \leq -\zeta, \\ \frac{1}{\mu} \frac{1}{|\zeta|}(x + 1 + 2/|\zeta|), & x \geq -\zeta. \end{cases}$$

Before proceeding to bound $|f''_h(x)|$ and $|f'''_h(x)|$, we first verify (C.4), or

$$\lim_{x \rightarrow \pm\infty} \nu(x) f''_h(x) = 0.$$

To do so, we rearrange the Poisson equation (3.3) to get

$$\nu(x) f''_h(x) = -\nu(x) \frac{b(x)}{\mu} f'_h(x) + \frac{\nu(x)}{\mu} [\mathbb{E}h(Y(\infty)) - h(x)].$$

It is now obvious that

$$\lim_{x \rightarrow \pm\infty} \nu(x) f''_h(x) \rightarrow 0 = 0,$$

because $h(x) \in \text{Lip}(1)$, the drift $b(x)$ is piecewise linear, and $f'_h(x)$ is bounded as in (C.18), but on the other hand $\nu(x)$ decays exponentially fast as $x \rightarrow \infty$, and decays even faster as $x \rightarrow -\infty$. To bound $|f''_h(x)|$, we use (C.7) and (C.8), together with the facts that $\|h'\| \leq 1$ and

$$b'(x) = -\mu 1(x < -\zeta), \quad x \in \mathbb{R},$$

to see that

$$(C.19) \quad |f''_h(x)| \leq \frac{1}{\nu(x)} \int_{-\infty}^x \frac{1}{\mu} (1 + \mu |f'_h(y)| 1(y < -\zeta)) \nu(y) dy$$

$$(C.20) \quad |f''_h(x)| \leq \frac{1}{\nu(x)} \int_x^{\infty} \frac{1}{\mu} (1 + \mu |f'_h(y)| 1(y < -\zeta)) \nu(y) dy.$$

We know $|f'_h(x)|$ is bounded as in (C.18). For $x \leq -\zeta$, we apply (C.10) to (C.19) and for $x \geq 0$ we apply (C.11) to (C.20) to conclude that

$$(C.21) \quad \mu |f''_h(x)| \leq \begin{cases} \sqrt{\frac{\pi}{2}} (1 + 6.5 + 4.2/|\zeta|), & x \leq 0, \\ \min \left\{ \sqrt{2\pi} e^{\frac{1}{2}\zeta^2}, \sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|} \right\} (1 + 6.5 + 4.2/|\zeta|), & x \in [0, -\zeta], \\ \frac{1}{|\zeta|}, & x \geq -\zeta. \end{cases}$$

By considering separately the cases when $|\zeta| \leq 1$ and $|\zeta| \geq 1$, we see that

$$(C.22) \quad \min \left\{ \sqrt{2\pi} e^{\frac{1}{2}\zeta^2}, \sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|} \right\} \leq 4.2,$$

and therefore,

$$(C.23) \quad |f_h''(x)| \leq \begin{cases} \frac{32}{\mu}(1 + 1/|\zeta|), & x \leq -\zeta, \\ \frac{1}{\mu|\zeta|}, & x \geq -\zeta. \end{cases}$$

Lastly, we bound $|f_h'''(x)|$, which exists for all $x \in \mathbb{R}$ where $h'(x)$ and $b'(x)$ exist. The fact that $h(x) \in \text{Lip}(1)$ together with (C.9) tells us that

$$|f_h'''(x)| \leq \frac{1}{\mu} (1 + |f_h''(x)b(x)| + |f_h'(x)b'(x)|).$$

For $x \geq -\zeta$, we use the forms of $b(x)$ and $b'(x)$ together with the bounds on $|f_h'(x)|$ and $|f_h''(x)|$ in (C.18) and (C.23) to see that

$$|f_h'''(x)| \leq \frac{1}{\mu} \left(1 + \mu |\zeta| \frac{1}{\mu |\zeta|} \right).$$

Although tempting, it is not sufficient to use the bound on $|f_h''(x)|$ in (C.21) and the form of $b(x)$ to bound $|f_h''(x)b(x)|$ for all $x \leq -\zeta$. Instead, we multiply both sides of (C.19) and (C.20) by $|b(x)|$ to see that

$$(C.24) \quad \begin{aligned} \|f_h''(x)b(x)\| &\leq \left(1 + \sup_{y \leq x} \mu |f_h'(y)| \right) \frac{|b(x)|}{\mu \nu(x)} \int_{-\infty}^x \nu(y) dy, \quad x \leq 0, \\ \|f_h''(x)b(x)\| &\leq \left(1 + \sup_{y \in [x, -\zeta]} \mu |f_h'(y)| \right) \frac{|b(x)|}{\mu \nu(x)} \int_x^{\infty} \nu(y) dy, \quad x \in [0, -\zeta]. \end{aligned}$$

By invoking (C.14) and (C.15) together with the bound on $|f_h'(x)|$ from (C.18), we conclude that

$$|f_h'''(x)| \leq \frac{1}{\mu} (1 + 2(1 + 6.5 + 4.2/|\zeta|) + 6.5 + 4.2/|\zeta|), \quad x \leq -\zeta.$$

Therefore, for those $x \in \mathbb{R}$ where $h'(x)$ and $b'(x)$ exist,

$$|f_h'''(x)| \leq \begin{cases} \frac{1}{\mu}(23 + 13/|\zeta|), & x \leq -\zeta, \\ \frac{2}{\mu}, & x \geq -\zeta. \end{cases}$$

This concludes the proof of Lemma 4. $\square$

PROOF OF LEMMA 12 . Recall that in the Erlang-C model, we set $\alpha = 0$, which makes

$$\zeta = \delta(x(\infty) - n) = \delta\left(\frac{\lambda}{\mu} - n\right) < 0, \quad b(x) = \begin{cases} -\mu x, & x \leq -\zeta, \\ \mu \zeta, & x \geq -\zeta. \end{cases}$$

The density of $Y(\infty)$ has the form

$$(C.25) \quad \nu(x) = \begin{cases} a_- e^{-\frac{1}{2}x^2}, & x \leq -\zeta, \\ a_+ e^{-|\zeta|x}, & x \geq -\zeta, \end{cases}$$

where a_- and a_+ are normalizing constants that make $\nu(x)$ continuous at the point $x = -\zeta$.

In the proof of this lemma we often rely on the fact that for any $c > 0$ and $x \geq 0$,

$$(C.26) \quad e^{\frac{1}{2}cx^2} \int_x^\infty e^{-\frac{1}{2}cy^2} dy \leq \int_0^\infty e^{-\frac{1}{2}cy^2} dy = \sqrt{\frac{\pi}{2c}}.$$

One can verify that the left hand side of (C.26) peaks at $x = 0$ by using the bound

$$(C.27) \quad \int_x^\infty e^{-\frac{1}{2}cy^2} dy \leq \int_x^\infty \frac{cy}{cx} e^{-\frac{1}{2}cy^2} dy = \frac{e^{-cx^2/2}}{cx}$$

to see that the derivative of the left side of (C.26) is negative for $x > 0$. We now prove (C.10). When $x \leq 0$, we use (C.26) and the symmetry of the function $e^{-\frac{1}{2}y^2}$ to see that

$$\frac{1}{\nu(x)} \int_{-\infty}^x \nu(y) dy = e^{\frac{1}{2}x^2} \int_{-\infty}^x e^{-\frac{1}{2}y^2} dy \leq \sqrt{\frac{\pi}{2}},$$

and when $x \in [0, -\zeta]$, we use (C.26) again to get

$$\begin{aligned} \frac{1}{\nu(x)} \int_{-\infty}^x \nu(y) dy &= e^{\frac{1}{2}x^2} \int_{-\infty}^0 e^{-\frac{1}{2}y^2} dy + e^{\frac{1}{2}x^2} \int_0^x e^{-\frac{1}{2}y^2} dy \leq 2e^{\frac{1}{2}\zeta^2} \int_{-\infty}^0 e^{-\frac{1}{2}y^2} dy \\ &= 2\sqrt{\frac{\pi}{2}} e^{\frac{1}{2}\zeta^2}. \end{aligned}$$

We now prove (C.11). Observe that when $x \geq -\zeta$,

$$\frac{1}{\nu(x)} \int_x^\infty \nu(y) dy = e^{|\zeta|x} \int_x^\infty e^{-|\zeta|y} dy = \frac{1}{|\zeta|},$$

and when $x \in [0, -\zeta]$, we use the fact that $a_- = a_+ e^{-\frac{1}{2}\zeta^2}$ together with (C.26) to see that

$$\begin{aligned} \frac{1}{\nu(x)} \int_x^\infty \nu(y) dy &= \frac{a_+}{a_-} e^{\frac{1}{2}x^2} \int_{-\zeta}^\infty e^{-|\zeta|y} dy + e^{\frac{1}{2}x^2} \int_x^{-\zeta} e^{-\frac{1}{2}y^2} dy \\ &\leq e^{\frac{1}{2}\zeta^2} e^{\frac{1}{2}x^2} \left(\frac{1}{|\zeta|} e^{-\zeta^2} \right) + \sqrt{\frac{\pi}{2}} \\ &\leq \frac{1}{|\zeta|} + \sqrt{\frac{\pi}{2}}. \end{aligned}$$

Moving on to show (C.12), when $x \leq 0$ we have

$$\frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy = e^{\frac{1}{2}x^2} \int_{-\infty}^x -ye^{-\frac{1}{2}y^2} dy = 1,$$

and when $x \in [0, -\zeta]$,

$$\begin{aligned} \frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy &= e^{\frac{1}{2}x^2} \int_{-\infty}^0 -ye^{-\frac{1}{2}y^2} dy + e^{\frac{1}{2}x^2} \int_0^x ye^{-\frac{1}{2}y^2} dy \\ &= e^{\frac{1}{2}x^2} + e^{\frac{1}{2}x^2} (1 - e^{-\frac{1}{2}x^2}). \end{aligned}$$

We now prove (C.13). When $x \in [0, -\zeta]$, we again use that $a_- = a_+ e^{-\frac{1}{2}\zeta^2}$ to obtain

$$\begin{aligned} \frac{1}{\nu(x)} \int_x^\infty |y| \nu(y) dy &= e^{\frac{1}{2}x^2} \int_x^{-\zeta} ye^{-\frac{1}{2}y^2} dy + \frac{a_+}{a_-} e^{\frac{1}{2}x^2} \int_{-\zeta}^\infty ye^{-|\zeta|y} dy \\ &= e^{\frac{1}{2}x^2} (e^{-\frac{1}{2}x^2} - e^{-\frac{1}{2}\zeta^2}) + e^{\frac{1}{2}\zeta^2} e^{\frac{1}{2}x^2} \left(1 + \frac{1}{\zeta^2}\right) e^{-\zeta^2} \\ &\leq 2 + \frac{1}{\zeta^2}. \end{aligned}$$

When $x \geq -\zeta$,

$$\frac{1}{\nu(x)} \int_x^\infty |y| \nu(y) dy = e^{|\zeta|x} \int_x^\infty ye^{-|\zeta|y} dy = \frac{x}{|\zeta|} + \frac{1}{\zeta^2}.$$

We move on to prove (C.14) and (C.15). When $x \leq 0$, we use (C.27) to see that

$$\frac{|b(x)|}{\mu\nu(x)} \int_{-\infty}^x \nu(y) dy = -xe^{\frac{x^2}{2}} \int_{-\infty}^x e^{-\frac{1}{2}y^2} dy \leq -xe^{\frac{x^2}{2}} \left(-\frac{1}{x} e^{-\frac{1}{2}x^2}\right) = 1.$$

When $x \in [0, -\zeta]$, we also use the fact that $a_+ = a_- e^{\frac{1}{2}\zeta^2}$ to get

$$\begin{aligned} \frac{|b(x)|}{\mu\nu(x)} \int_x^\infty \nu(y) dy &= xe^{\frac{x^2}{2}} \int_x^{-\zeta} e^{-\frac{1}{2}y^2} dy + \frac{a_+}{a_-} xe^{\frac{x^2}{2}} \int_{-\zeta}^\infty e^{-|\zeta|y} dy \\ &\leq xe^{\frac{x^2}{2}} \left(\frac{1}{x} e^{-\frac{1}{2}x^2}\right) + e^{\frac{1}{2}\zeta^2} xe^{\frac{x^2}{2}} \left(\frac{1}{|\zeta|} e^{-\zeta^2}\right) \\ &= 1 + \frac{x}{|\zeta|} e^{\frac{1}{2}(x^2 - \zeta^2)} \leq 2. \end{aligned}$$

When $x \geq -\zeta$,

$$\frac{|b(x)|}{\mu\nu(x)} \int_x^\infty \nu(y) dy = |\zeta| e^{|\zeta|x} \int_x^\infty e^{-|\zeta|y} dy = 1.$$

Lastly we prove (C.16). Letting $V(x) = x^2$, and recalling the form of G_Y from (3.2), we consider

$$(C.28) \quad G_Y V(x) = 2x\mu(\zeta + (x + \zeta)^-) + 2\mu = -2\mu x^2 1(x < -\zeta) - 2x\mu |\zeta| 1(x \geq -\zeta) + 2\mu.$$

By the standard Foster-Lyapunov condition (see for example [13, Theorem 4.3]), this implies that

$$2\mathbb{E}[(Y(\infty))^2 1(Y(\infty) < -\zeta)] + 2|\zeta| \mathbb{E}[Y(\infty) 1(Y(\infty) \geq -\zeta)] \leq 2,$$

and in particular,

$$\begin{aligned} \mathbb{E}[Y(\infty) 1(Y(\infty) \geq -\zeta)] &\leq \frac{1}{|\zeta|}, \\ \mathbb{E}[Y(\infty) 1(Y(\infty) < -\zeta)] &\leq \sqrt{\mathbb{E}[(Y(\infty))^2 1(Y(\infty) < -\zeta)]} \leq 1, \end{aligned}$$

where we applied Jensen's inequality in the second set of inequalities. This concludes the proof of Lemma 12. $\square$

C.1.2. Proof Outline for Lemma 5: The Underloaded System. In this section we prove the underloaded case of Lemma 5. The proof is very similar to that of Lemma 4. Therefore, we only outline some key intermediate steps needed to obtain the results. We will need the following version of Lemma 12.

LEMMA 13. *Consider the Erlang-A model ($\alpha > 0$) with $0 < R \leq n$, and let $\nu(x)$ be the density of $Y(\infty)$. Then*

$$(C.29) \quad \frac{1}{\nu(x)} \int_{-\infty}^x \nu(y) dy \leq \begin{cases} \sqrt{\frac{\pi}{2}}, & x \leq 0, \\ \sqrt{2\pi} e^{\frac{\zeta^2}{2}}, & x \in [0, -\zeta], \end{cases}$$

$$(C.30) \quad \frac{1}{\nu(x)} \int_x^{\infty} \nu(y) dy \leq \begin{cases} \sqrt{\frac{\pi}{2}} + \sqrt{\frac{\pi}{2}} \frac{\mu}{\alpha} \wedge \frac{1}{|\zeta|}, & x \in [0, -\zeta], \\ \sqrt{\frac{\pi}{2}} \frac{\mu}{\alpha} \wedge \frac{1}{|\zeta|}, & x \geq -\zeta, \end{cases}$$

$$(C.31) \quad \frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy \leq \begin{cases} 1, & x \leq 0, \\ 2e^{\frac{\zeta^2}{2}} - 1, & x \in [0, -\zeta], \end{cases}$$

$$(C.32) \quad \frac{1}{\nu(x)} \int_x^{\infty} |y| \nu(y) dy \leq \begin{cases} 2 + \frac{1}{\zeta^2}, & x \in [0, -\zeta], \\ 1 + \frac{\mu}{\alpha}, & x \geq -\zeta, \end{cases}$$

$$(C.33) \quad \frac{|b(x)|}{\mu \nu(x)} \int_{-\infty}^x \nu(y) dy \leq 1, \quad x \leq 0,$$

$$(C.34) \quad \frac{|b(x)|}{\mu \nu(x)} \int_x^{\infty} \nu(y) dy \leq 2, \quad x \geq 0,$$

$$(C.35) \quad \mathbb{E} |Y(\infty)| \leq 1 + \sqrt{\frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|}.$$

To prove this lemma, we first observe that

$$b(x) = \begin{cases} -\mu x, & x \leq -\zeta, \\ -\alpha(x + \zeta) + \mu\zeta, & x \geq -\zeta, \end{cases}$$

and

$$(C.36) \quad \nu(x) = \begin{cases} a_- e^{-\frac{1}{2}x^2}, & x \leq -\zeta, \\ a_+ e^{-\frac{\alpha}{2\mu}(x+\zeta-\frac{\mu}{\alpha}\zeta)^2}, & x \geq -\zeta, \end{cases}$$

where a_- and a_+ are normalizing constants that make $\nu(x)$ continuous and integrate to one. By comparing the density in (C.36) to (C.25) for the region $x \leq -\zeta$, we immediately see that (C.29), (C.31) and (C.33) are restatements of (C.10), (C.12), and (C.14) from Lemma 12, and hence have already been established. The proof of (C.35) involves applying G_Y to the Lyapunov function $V(x) = x^2$ to see that

$$\begin{aligned} G_Y V(x) &= -2\mu x^2 1(x < -\zeta) + 2(-\alpha x^2 + x\zeta(\mu - \alpha)) 1(x \geq -\zeta) + 2\mu \\ &\leq -2\mu x^2 1(x < -\zeta) - 2(\alpha \wedge \mu) x^2 1(x \geq -\zeta) + 2\mu, \end{aligned}$$

and

$$\begin{aligned} G_Y V(x) &= -2\mu x^2 1(x < -\zeta) + 2(-\alpha x(x + \zeta) - \mu |\zeta| x) 1(x \geq -\zeta) + 2\mu \\ &\leq -2\mu x^2 1(x < -\zeta) - 2\mu |\zeta| x 1(x \geq -\zeta) + 2\mu. \end{aligned}$$

One can compare these inequalities to (C.28) in the proof of Lemma 12 to see that (C.35) follows by the Foster-Lyapunov condition.

We now go over the proofs of (C.30), (C.32) and (C.34). We first prove (C.30) when $x \in [0, -\zeta]$. Observe that

$$\begin{aligned} \frac{1}{\nu(x)} \int_x^\infty \nu(y) dy &= e^{\frac{1}{2}x^2} \int_x^{|\zeta|} e^{-\frac{1}{2}y^2} dy + \frac{a_+}{a_-} e^{\frac{1}{2}x^2} \int_{|\zeta|}^\infty e^{-\frac{\alpha}{2\mu}(y+\zeta-\frac{\mu}{\alpha}\zeta)^2} dy \\ &\leq e^{\frac{1}{2}x^2} \int_x^\infty e^{-\frac{1}{2}y^2} dy + e^{\frac{1}{2}(x^2-\zeta^2)} e^{\frac{\alpha}{2\mu}(\frac{\mu}{\alpha}\zeta)^2} \int_{\frac{\mu}{\alpha}|\zeta|}^\infty e^{-\frac{\alpha}{2\mu}y^2} dy \\ (C.37) \quad &\leq \sqrt{\frac{\pi}{2}} + \sqrt{\frac{\pi}{2} \frac{\mu}{\alpha}} \wedge \frac{1}{|\zeta|} \end{aligned}$$

where in the first inequality we used a change of variables and the fact that $a_+/a_- = e^{-\zeta^2/2} e^{\frac{\alpha}{2\mu}(\frac{\mu}{\alpha}\zeta)^2}$, and in the last inequality we used both (C.26) and (C.27), and the fact

that $e^{\frac{1}{2}(x^2-\zeta^2)} \leq 1$. The rest of (C.30) is proved identically. We now prove (C.32). When $x \in [0, -\zeta]$,

$$\begin{aligned} \frac{1}{\nu(x)} \int_x^\infty |y| \nu(y) dy &= e^{\frac{1}{2}x^2} \int_x^{-\zeta} y e^{-\frac{1}{2}y^2} dy + \frac{a_+}{a_-} e^{\frac{1}{2}x^2} \int_{-\zeta}^\infty y e^{-\frac{\alpha}{2\mu}(y+\zeta-\frac{\mu}{\alpha}\zeta)^2} dy \\ &= (1 - e^{\frac{1}{2}(x^2-\zeta^2)}) + e^{\frac{\alpha}{2\mu}(\frac{\mu}{\alpha}\zeta)^2} e^{\frac{1}{2}(x^2-\zeta^2)} \int_{-\zeta}^\infty y e^{-\frac{\alpha}{2\mu}(y+\zeta-\frac{\mu}{\alpha}\zeta)^2} dy \\ &\leq 1 + e^{\frac{\alpha}{2\mu}(\frac{\mu}{\alpha}\zeta)^2} \int_{-\zeta}^\infty y e^{-\frac{\alpha}{2\mu}((y+\zeta)^2 - 2\frac{\mu}{\alpha}(y+\zeta)\zeta + (\frac{\mu}{\alpha}\zeta)^2)} dy \\ &\leq 1 + \int_{-\zeta}^\infty y e^{(y+\zeta)\zeta} dy = 1 + 1 + \frac{1}{\zeta^2}, \end{aligned}$$

and when $x \geq -\zeta$,

$$\begin{aligned} \frac{1}{\nu(x)} \int_x^\infty |y| \nu(y) dy &= e^{\frac{\alpha}{2\mu}(x+\zeta-\frac{\mu}{\alpha}\zeta)^2} \int_x^\infty y e^{-\frac{\alpha}{2\mu}(y+\zeta-\frac{\mu}{\alpha}\zeta)^2} dy \\ &= e^{\frac{\alpha}{2\mu}(x+\zeta-\frac{\mu}{\alpha}\zeta)^2} \int_{x+\zeta-\frac{\mu}{\alpha}\zeta}^\infty y e^{-\frac{\alpha}{2\mu}y^2} dy \\ &\quad + (1 - \mu/\alpha) |\zeta| e^{\frac{\alpha}{2\mu}(x+\zeta-\frac{\mu}{\alpha}\zeta)^2} \int_{x+\zeta-\frac{\mu}{\alpha}\zeta}^\infty e^{-\frac{\alpha}{2\mu}y^2} dy \\ &\leq \frac{\mu}{\alpha} + |\zeta| \frac{1}{\frac{\alpha}{\mu}(x+\zeta-\frac{\mu}{\alpha}\zeta)} \leq \frac{\mu}{\alpha} + 1, \end{aligned}$$

where in the last inequality we used (C.27). Lastly, the argument for (C.34) is very similar to the chain of inequalities in (C.37), and we leave the details to the reader.

We now describe how to prove Lemma 5. To prove (3.37), we apply Lemma 13 to (C.5) and (C.6) just like in (C.17) of Lemma 4. Using these bounds on $f'_h(x)$, we argue (C.4) just like in the proof of Lemma 4. We now describe how to prove (3.38). When $x \leq 0$, we apply (3.37) and (C.29) to (C.7), and when $x \geq -\zeta$ we apply (3.37) and (C.30) to (C.8). The last region, when $x \in [0, -\zeta]$, has to be handled differently depending on the size of $|\zeta|$. When $|\zeta| \leq 1$, we just apply (3.37) and (C.29) to (C.7). However, when $|\zeta| \geq 1$, we manipulate (C.8) to see that

$$(C.38) \quad f''_h(x) = -\frac{1}{\nu(x)} \int_x^{-\zeta} \frac{1}{\mu} (-h'(y) + \mu f'_h(y)) \nu(y) dy$$

$$(C.39) \quad -\frac{\nu(-\zeta)}{\nu(x)} \frac{1}{\nu(-\zeta)} \int_{-\zeta}^\infty \frac{1}{\mu} (-h'(y) + \alpha f'_h(y)) \nu(y) dy.$$

We then apply (3.37), (C.30), and the fact that $\nu(-\zeta)/\nu(x) \leq 1$ to conclude (3.38). The proof of (3.40) relies on (C.9), which tells us that

$$|f'''_h(x)| \leq \frac{1}{\mu} [1 + |f''_h(x)b(x)| + |f'_h(x)b'(x)|].$$

Bounding $|f_h'(x)b'(x)|$ is simple, and only relies on (3.37). The term $|f_h''(x)b(x)|$ is bounded similarly to the way it is done in Lemma 4; see for instance (C.24). Namely, for $x \leq 0$ we multiply both sides of (C.7) by $b(x)$ and apply (3.37) with (C.33), and when $x \geq -\zeta$ we multiply both sides of (C.8) by $b(x)$ and apply (3.37) with (C.34). Lastly, when $x \in [0, -\zeta]$, we manipulate (C.8) to get

$$\begin{aligned} b(x)f_h''(x) = & -\frac{b(x)}{\nu(x)} \int_x^{-\zeta} \frac{1}{\mu} (-h'(y) + \mu f_h'(y)) \nu(y) dy \\ & -\frac{b(x)\nu(-\zeta)}{\nu(x)b(-\zeta)} \frac{b(-\zeta)}{\nu(-\zeta)} \int_{-\zeta}^{\infty} \frac{1}{\mu} (-h'(y) + \alpha f_h'(y)) \nu(y) dy. \end{aligned}$$

We then apply (3.37), (C.34), and the fact that $|\frac{b(x)\nu(-\zeta)}{\nu(x)b(-\zeta)}| \leq 1$ to get the required bounds on $|b(x)f_h''(x)|$ and conclude (3.40). This concludes the proof outline for Lemma 5.

C.1.3. Proof Outline for Lemma 5: The Overloaded System. In this section we prove the overloaded case of Lemma 5. The proof is similar to that of Lemma 4. Therefore, we only outline some key intermediate steps needed to obtain the results. We will need the following version of Lemma 12.

LEMMA 14. *Consider the Erlang-A model ($\alpha > 0$) with $R \geq n$, and let $\nu(x)$ be the density of $Y(\infty)$. Then*

$$(C.40) \quad \frac{1}{\nu(x)} \int_{-\infty}^x \nu(y) dy \leq \begin{cases} \sqrt{\frac{\pi}{2}} \wedge \frac{\mu}{\alpha\zeta}, & x \leq -\zeta, \\ \sqrt{\frac{\pi}{2}} + \sqrt{\frac{\pi}{2}} \frac{\mu}{\alpha} \wedge \zeta, & x \in [-\zeta, 0], \end{cases}$$

$$(C.41) \quad \frac{1}{\nu(x)} \int_x^{\infty} \nu(y) dy \leq \begin{cases} \sqrt{2\pi} \frac{\mu}{\alpha} e^{\frac{\alpha}{2\mu}\zeta^2}, & x \in [-\zeta, 0], \\ \sqrt{\frac{\pi}{2}} \frac{\mu}{\alpha}, & x \geq 0, \end{cases}$$

$$(C.42) \quad \frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy \leq \begin{cases} 1 + \sqrt{\frac{\pi}{2}} \zeta \wedge \frac{\mu}{\alpha}, & x \leq -\zeta, \\ \frac{\mu}{\alpha} + 1, & x \in [-\zeta, 0], \end{cases}$$

$$(C.43) \quad \frac{1}{\nu(x)} \int_x^{\infty} |y| \nu(y) dy \leq \begin{cases} 2\frac{\mu}{\alpha} e^{\frac{\alpha}{2\mu}\zeta^2}, & x \in [-\zeta, 0], \\ \frac{\mu}{\alpha}, & x \geq 0, \end{cases}$$

$$(C.44) \quad \frac{|b(x)|}{\mu\nu(x)} \int_{-\infty}^x \nu(y) dy \leq 2, \quad x \leq 0,$$

$$(C.45) \quad \frac{|b(x)|}{\mu\nu(x)} \int_x^{\infty} \nu(y) dy \leq 1, \quad x \geq 0,$$

$$(C.46) \quad \mathbb{E} |Y(\infty)| \leq \sqrt{\frac{\mu}{\alpha}} + 1.$$

To prove this lemma, we first observe that

$$b(x) = \begin{cases} -\mu(x + \zeta) + \alpha\zeta, & x \leq -\zeta, \\ -\alpha x, & x \geq -\zeta, \end{cases}$$

and

$$(C.47) \quad \nu(x) = \begin{cases} a_- e^{-\frac{1}{2}\left(x+\zeta-\frac{\alpha}{\mu}\zeta\right)^2}, & x \leq -\zeta, \\ a_+ e^{-\frac{\alpha}{2\mu}x^2}, & x \geq -\zeta, \end{cases}$$

where a_- and a_+ are normalizing constants that make $\nu(x)$ continuous and integrate to one. Observe that in the region $x \geq -\zeta$, the density in (C.47) looks very similar to the density in (C.25) in the region $x \leq -\zeta$. Hence, one can check that the arguments needed to prove Lemma 14's (C.41), (C.43), and (C.45) are nearly identical to the arguments used to prove Lemma 12's (C.10), (C.12), and (C.14), respectively.

The proof of (C.46) involves applying G_Y to the Lyapunov function $V(x) = x^2$ to see that

$$\begin{aligned} G_Y V(x) &= -2\alpha x^2 1(x > -\zeta) + 2(-\mu x^2 + x\zeta(\alpha - \mu)) 1(x \leq -\zeta) + 2\mu \\ &\leq -2\alpha x^2 1(x > -\zeta) - 2(\alpha \wedge \mu) x^2 1(x \leq -\zeta) + 2\mu. \end{aligned}$$

One can compare this inequality to (C.28) in the proof of Lemma 12 to see that (C.46) follows by the Foster-Lyapunov condition.

We now describe how to prove (C.40), (C.42), and (C.44). To prove (C.40) we use a series of arguments similar to those in (C.37), where we proved (C.30) of Lemma 13. We now prove (C.42). When $x \leq -\zeta$,

$$\begin{aligned} \frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy &= e^{\frac{1}{2}(x+\zeta-\frac{\alpha}{\mu}\zeta)^2} \int_{-\infty}^x -y e^{-\frac{1}{2}(y+\zeta-\frac{\alpha}{\mu}\zeta)^2} dy \\ &= e^{\frac{1}{2}(x+\zeta-\frac{\alpha}{\mu}\zeta)^2} \int_{-\infty}^{x+\zeta-\frac{\alpha}{\mu}\zeta} -y e^{-\frac{1}{2}y^2} dy \\ &\quad + (1 - \alpha/\mu)\zeta e^{\frac{1}{2}(x+\zeta-\frac{\alpha}{\mu}\zeta)^2} \int_{-\infty}^{x+\zeta-\frac{\alpha}{\mu}\zeta} e^{-\frac{1}{2}y^2} dy \\ (C.48) \quad &\leq 1 + \zeta \left(\sqrt{\frac{\pi}{2}} \wedge \frac{1}{\frac{\alpha}{\mu}\zeta - x - \zeta} \right) \leq 1 + \sqrt{\frac{\pi}{2}} \zeta \wedge \frac{\mu}{\alpha}, \end{aligned}$$

where in the last inequality we used both (C.26) and (C.27). When $x \in [-\zeta, 0]$,

$$\frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy = \frac{a_-}{a_+} e^{\frac{\alpha}{2\mu}x^2} \int_{-\infty}^{-\zeta} -y e^{-\frac{1}{2}(y+\zeta-\frac{\alpha}{\mu}\zeta)^2} dy + e^{\frac{\alpha}{2\mu}x^2} \int_{-\zeta}^x -y e^{\frac{\alpha}{2\mu}y^2} dy.$$

Repeating arguments from (C.48) and using the fact that $a_-/a_+ = e^{-\frac{\alpha}{2\mu}\zeta^2} e^{\frac{1}{2}(\frac{\alpha}{\mu}\zeta)^2}$, we can show that the first term above satisfies

$$\frac{a_-}{a_+} e^{\frac{\alpha}{2\mu}x^2} \int_{-\infty}^{-\zeta} -y e^{-\frac{1}{2}(y+\zeta-\frac{\alpha}{\mu}\zeta)^2} dy \leq e^{\frac{\alpha}{2\mu}(x^2-\zeta^2)} \left(1 + \frac{\mu}{\alpha}\right),$$

and by computing the second term explicitly, we conclude that

$$\begin{aligned} \frac{1}{\nu(x)} \int_{-\infty}^x |y| \nu(y) dy &\leq e^{\frac{\alpha}{2\mu}(x^2-\zeta^2)} \left(1 + \frac{\mu}{\alpha}\right) + \frac{\mu}{\alpha} (1 - e^{\frac{\alpha}{2\mu}(x^2-\zeta^2)}) \\ &\leq 1 + \frac{\mu}{\alpha}, \end{aligned}$$

which proves (C.42). Lastly, it is not hard to see that (C.44) follows from a straightforward application of (C.27).

Having argued Lemma 14, we now use it to prove the bounds in (3.41)–(3.46). To prove (3.41), we apply Lemma 14 to (C.5) and (C.6) just like in (C.17) of Lemma 4. Using these bounds on $f'_h(x)$, we argue (C.4) just like in the proof of Lemma 4. We now describe how to prove (3.42). When $x \leq -\zeta$, we apply (3.41) and (C.40) to (C.7). When $x \geq -\zeta$, instead of using the expressions for $f''_h(x)$ in (C.7) and (C.8) like we would usually do, we instead apply (3.41) to the bound

$$|f''_h(x)| \leq \frac{1}{\mu} |f'_h(x)| |b(x)| + \frac{1}{\mu} (|x| + \mathbb{E}|Y(\infty)|), \quad x \in \mathbb{R},$$

which follows by rewriting the Poisson equation (3.3) and using the Lipschitz property of $h(x)$. We now prove (3.44)–(3.46). We recall (C.9) to see that

$$|f'''_h(x)| \leq \frac{1}{\mu} [1 + |f''_h(x)b(x)| + |f'_h(x)b'(x)|].$$

Bounding $|f'_h(x)b'(x)|$ is simple, and only relies on (3.41). The other term, $|f''_h(x)b(x)|$, is bounded as follows. To prove (3.44), i.e. when $x \leq -\zeta$, we multiply both sides of (C.7) by $b(x)$, and apply (3.41) and (C.44) to the result. When $x \geq -\zeta$ then

$$|f''_h(x)b(x)| = \alpha |x| |f''_h(x)|,$$

and the difference between (3.45) and (3.46) lies in the way that the quantity above is bounded. To get (3.45), we simply apply the bounds on $f''_h(x)$ from (3.42) to the right hand side above.

To prove (3.46), we will first argue that

$$(C.49) \quad |f'''_h(x)| \leq \begin{cases} \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) + \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right)^2 |x|, & x \in [-\zeta, 0], \\ \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right), & x \geq 0, \end{cases}$$

where C is some positive constant independent of everything else; this will imply (3.46). The only difference between the proof of (C.49) and the bound on $f_h'''(x)$ in (3.45) is in how $|f_h''(x)b(x)|$ is bounded; we now describe the different way to bound $|f_h''(x)b(x)|$. When $x \geq 0$, we multiply both sides of (C.8) by $b(x)$ and use the bounds in (3.41) and (C.45) to bound $|f_h''(x)b(x)|$. When $x \in [-\zeta, 0]$, we want to prove that

$$(C.50) \quad |f_h''(x)| \leq \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \left(1 + \sqrt{\frac{\mu}{\alpha}} \right) + \frac{C}{\mu} \left(\zeta \wedge \frac{\mu^2}{\alpha^2 \zeta} \right),$$

which, after considering separately the cases when $\zeta \leq \mu/\alpha$ and $\zeta \geq \mu/\alpha$, implies that

$$|f_h''(x)| \leq \frac{C}{\mu} \left(\frac{\alpha}{\mu} + \sqrt{\frac{\alpha}{\mu}} + 1 \right) \left(1 + \sqrt{\frac{\mu}{\alpha}} \right) + \frac{C}{\alpha}.$$

We can then use this fact to bound $|f_h''(x)b(x)| = \alpha|x||f_h''(x)|$. To prove (C.50) for $\zeta \leq \sqrt{\mu/\alpha}$, we bound (C.8) using (3.41) and (C.41). To prove (C.50) for $\zeta \geq \sqrt{\mu/\alpha}$, we bound (C.7) using (3.41) and (C.40). We point out that to bound (C.7) we need to perform a manipulation similar to the one in (C.39). This concludes the proof outline for the overloaded case.

C.2. Kolmogorov Gradient Bounds: Proof of Lemmas 6 and 7. For the remainder of this section, we take $\mathcal{H} = \mathcal{H}_K$ in (3.3). With this choice of test functions, any solution to the Poisson equation will have a discontinuity in its second derivative, which makes the gradient bounds for it differ from the Wasserstein setting. Fix $a \in \mathbb{R}$ and consider the Poisson equation

$$G_Y f_a(x) = \mu f_a''(x) + b(x) f_a'(x) = F_Y(a) - 1_{(-\infty, a]}(x),$$

where $F_Y(x)$ is the distribution function of $Y(\infty)$. If $f_a(x)$ is a solution the Poisson equation with $a_2 = 0$, then just as in (C.1) and (C.2),

$$\begin{aligned} f_a'(x) &= \frac{1}{\mu\nu(x)} \int_{-\infty}^x (F_Y(a) - 1_{(-\infty, a]}(y)) \nu(y) dy, \\ f_a'(x) &= -\frac{1}{\mu\nu(x)} \int_x^{\infty} (F_Y(a) - 1_{(-\infty, a]}(y)) \nu(y) dy, \end{aligned}$$

which immediately implies that

$$|f_a'(x)| \leq \frac{1}{\mu\nu(x)} \min \left\{ \int_{-\infty}^x \nu(y) dy, \int_x^{\infty} \nu(y) dy \right\}.$$

Furthermore,

$$f_a''(x) = \frac{1}{\mu} [F_Y(a) - 1_{(-\infty, a]}(x) - b(x) f_a'(x)].$$

We now prove the Kolmogorov gradient bounds for the Erlang-C model.

PROOF OF LEMMA 6. First of all, by (C.10) and (C.11),

$$(C.51) \quad \mu |f'_a(x)| \leq \begin{cases} \sqrt{\frac{\pi}{2}}, & x \leq 0, \\ \min \left\{ \sqrt{2\pi} e^{\frac{1}{2}\zeta^2}, \sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|} \right\}, & x \in [0, -\zeta], \\ \frac{1}{|\zeta|}, & x \geq -\zeta, \end{cases}$$

and (C.22) implies that

$$\min \left\{ \sqrt{2\pi} e^{\frac{1}{2}\zeta^2}, \sqrt{\frac{\pi}{2}} + \frac{1}{|\zeta|} \right\} \leq 5,$$

which proves the bounds for $f'_a(x)$. Second, (C.14) and (C.15) imply that for all $x \in \mathbb{R}$,

$$(C.52) \quad |f''_a(x)| \leq \frac{1}{\mu} \left[1 + \frac{|b(x)|}{\mu\nu(x)} \min \left\{ \int_{-\infty}^x \nu(y) dy, \int_x^{\infty} \nu(y) dy \right\} \right] \leq 3/\mu,$$

where $f''_a(x)$ is understood to be the left derivative at the point $x = a$.

□

PROOF OF LEMMA 7. The proof of this lemma is almost identical to the proof of Lemma 6. By using the analogues of (C.14) and (C.15) from Lemmas 13 and 14, it's not hard to check that (C.52) holds for the Erlang-A model as well. To prove the bounds on $f'_a(x)$, we obtain inequalities similar to (C.51) by using analogues of (C.10) and (C.11) from Lemmas 13 and 14. These inequalities will imply (5.3) and (5.4) once we consider in them separately the cases when $|\zeta| \leq 1$ and $|\zeta| \geq 1$.

□

APPENDIX D: KOLMOGOROV PROOFS

In the section we prove Lemmas 8 and 9.

D.1. Proof of Lemma 8.

PROOF OF LEMMA 8. We let $F_W(w)$ and $F_{\tilde{X}}(x)$ be the distribution functions of W and $\tilde{X}(\infty)$, respectively. For any $a \in \mathbb{R}$, let $\tilde{a} = \delta(a - x(\infty))$. We want to show that

$$(D.1) \quad \begin{aligned} \mathbb{P}(\tilde{a} - \delta < \tilde{X}(\infty) \leq \tilde{a} + \delta) &= F_{\tilde{X}}(\tilde{a} + \delta) - F_{\tilde{X}}(\tilde{a} - \delta) \\ &\leq 2\delta\omega(F_W) + d_K(\tilde{X}(\infty), W) + 9\delta^2 + 8\delta^4. \end{aligned}$$

Define

$$k^* = \inf\{k \geq 0 : \nu_k \geq \nu_j, \text{ for all } j \neq k\}.$$

Then for any $\tilde{a} \in \mathbb{R}$,

$$F_{\tilde{X}}(\tilde{a} + \delta) - F_{\tilde{X}}(\tilde{a} - \delta) \leq 2\nu_{k^*},$$

because $\tilde{X}(\infty)$ takes at most two values in the interval $(\tilde{a} - \delta, \tilde{a} + \delta]$. Observe that by the flow balance equations, we know that for any $k \in \mathbb{Z}_+$,

$$\nu_k = \frac{d(k+1)}{\lambda} \nu_{k+1}.$$

Since k^* is the maximizer of $\{\nu_k\}$, we know that

$$d(k^*) \leq \lambda \leq d(k^* + 1) \leq \lambda + \mu,$$

where in the last inequality we have used the fact that the increase in departure rate between state k^* and $k^* + 1$ is at most μ . Likewise, $d(k^* + i) \leq \lambda + i\mu$ for $i = 2, 3$. Hence,

$$\begin{aligned} \nu_{k^*} &= \frac{d(k^* + 1)}{\lambda} \nu_{k^*+1} \leq \left(1 + \frac{\mu}{\lambda}\right) \nu_{k^*+1} \leq \nu_{k^*+1} + \delta^2, \\ \nu_{k^*} &= \frac{d(k^* + 1)}{\lambda} \frac{d(k^* + 2)}{\lambda} \nu_{k^*+2} \leq (1 + \delta^2)(1 + 2\delta^2) \nu_{k^*+2} \leq \nu_{k^*+2} + 3\delta^2 + 2\delta^4, \\ \nu_{k^*+1} &= \frac{d(k^* + 2)}{\lambda} \frac{d(k^* + 3)}{\lambda} \nu_{k^*+3} \leq (1 + 2\delta^2)(1 + 3\delta^2) \nu_{k^*+3} \leq \nu_{k^*+3} + 5\delta^2 + 6\delta^4, \end{aligned}$$

which implies that for any $\tilde{a} \in \mathbb{R}$,

$$\begin{aligned} F_{\tilde{X}}(\tilde{a} + \delta) - F_{\tilde{X}}(\tilde{a} - \delta) &\leq 2\nu_{k^*} \leq \nu_{k^*} + \nu_{k^*+1} + \delta^2 \\ &= F_{\tilde{X}}(\tilde{k}^* + \delta) - F_{\tilde{X}}(\tilde{k}^* - \delta) + \delta^2. \end{aligned}$$

There are now 4 cases to consider, with the first three being simple to handle. Recall that $\omega(F_W)$ is the modulus of continuity of $F_W(w)$.

1. If $F_W(\tilde{k}^* - \delta) \leq F_{\tilde{X}}(\tilde{k}^* - \delta)$ and $F_W(\tilde{k}^* + \delta) \geq F_{\tilde{X}}(\tilde{k}^* + \delta)$, then

$$(D.2) \quad F_{\tilde{X}}(\tilde{k}^* + \delta) - F_{\tilde{X}}(\tilde{k}^* - \delta) \leq F_W(\tilde{k}^* + \delta) - F_W(\tilde{k}^* - \delta) \leq 2\delta\omega(F_W).$$

2. If $F_W(\tilde{k}^* - \delta) \leq F_{\tilde{X}}(\tilde{k}^* - \delta)$ but $F_W(\tilde{k}^* + \delta) < F_{\tilde{X}}(\tilde{k}^* + \delta)$, then

$$\begin{aligned} F_{\tilde{X}}(\tilde{k}^* + \delta) - F_{\tilde{X}}(\tilde{k}^* - \delta) &\leq F_{\tilde{X}}(\tilde{k}^* + \delta) - F_W(\tilde{k}^* + \delta) + F_W(\tilde{k}^* + \delta) - F_W(\tilde{k}^* - \delta) \\ (D.3) \quad &\leq 2\delta\omega(F_W) + d_K(\tilde{X}(\infty), W). \end{aligned}$$

3. Similarly, if $F_W(\tilde{k}^* - \delta) > F_{\tilde{X}}(\tilde{k}^* - \delta)$ and $F_W(\tilde{k}^* + \delta) \geq F_{\tilde{X}}(\tilde{k}^* + \delta)$, then

$$\begin{aligned} F_{\tilde{X}}(\tilde{k}^* + \delta) - F_{\tilde{X}}(\tilde{k}^* - \delta) &\leq F_W(\tilde{k}^* + \delta) - F_W(\tilde{k}^* - \delta) + F_W(\tilde{k}^* - \delta) - F_{\tilde{X}}(\tilde{k}^* - \delta) \\ (D.4) \quad &\leq 2\delta\omega(F_W) + d_K(\tilde{X}(\infty), W). \end{aligned}$$

4. Suppose $F_W(\tilde{k}^* - \delta) > F_{\tilde{X}}(\tilde{k}^* - \delta)$ and $F_W(\tilde{k}^* + \delta) < F_{\tilde{X}}(\tilde{k}^* + \delta)$, then we need to use a different approach. We know that

$$\begin{aligned} F_{\tilde{X}}(\tilde{k}^* + \delta) - F_{\tilde{X}}(\tilde{k}^* - \delta) &= \nu_{k^*} + \nu_{k^*+1} \\ &\leq \nu_{k^*+2} + \nu_{k^*+3} + 8\delta^2 + 8\delta^4 \\ &= F_{\tilde{X}}(\tilde{k}^* + 3\delta) - F_{\tilde{X}}(\tilde{k}^* + \delta) + 8\delta^2 + 8\delta^4. \end{aligned}$$

Since $F_W(\tilde{k}^* + \delta) \leq F_{\tilde{X}}(\tilde{k}^* + \delta)$, we are either in case 1 or 2 for $F_{\tilde{X}}(\tilde{k}^* + 3\delta) - F_{\tilde{X}}(\tilde{k}^* + \delta)$, and hence we have

$$F_{\tilde{X}}(\tilde{k}^* + 3\delta) - F_{\tilde{X}}(\tilde{k}^* + \delta) \leq 2\delta\omega(F_W) + d_K(\tilde{X}(\infty), W).$$

This proves (D.1), concluding the proof of this lemma. □

D.2. Proof of Lemma 9 .

PROOF OF LEMMA 9. In the Erlang-C model,

$$\nu(x) = \begin{cases} a_- e^{-\frac{1}{2}x^2}, & x \leq -\zeta, \\ a_+ e^{-|\zeta|x}, & x \geq -\zeta. \end{cases}$$

To bound this density, we need to bound a_- and a_+ . We know that $\nu(x)$ must integrate to one, which implies that

$$a_- \int_{-\infty}^{-\zeta} e^{-\frac{1}{2}y^2} dy + a_+ \int_{-\zeta}^{\infty} e^{-|\zeta|y} dy = 1$$

Furthermore, since $\nu(x)$ is continuous at $x = -\zeta$,

$$a_- e^{-\frac{1}{2}\zeta^2} = a_+ e^{-\zeta^2}.$$

Combining these two facts, we see that

$$a_- = \frac{1}{\int_{-\infty}^{-\zeta} e^{-\frac{1}{2}y^2} dy + e^{\frac{1}{2}\zeta^2} \int_{-\zeta}^{\infty} e^{-|\zeta|y} dy} \leq \frac{1}{\int_{-\infty}^0 e^{-\frac{1}{2}y^2} dy} = \sqrt{\frac{2}{\pi}},$$

and

$$a_+ = \frac{1}{e^{-\frac{1}{2}\zeta^2} \int_{-\infty}^{-\zeta} e^{-\frac{1}{2}y^2} dy + \int_{-\zeta}^{\infty} e^{-|\zeta|y} dy} \leq \frac{1}{e^{-\frac{1}{2}\zeta^2} \int_{-\infty}^0 e^{-\frac{1}{2}y^2} dy} = e^{\frac{1}{2}\zeta^2} \sqrt{\frac{2}{\pi}}.$$

Therefore, for $x \leq -\zeta$,

$$|\nu(x)| \leq a_- \leq \sqrt{\frac{2}{\pi}},$$

and for $x \geq -\zeta$, we recall that $\zeta < 0$ to see that

$$|\nu(x)| \leq a_+ e^{-|\zeta|x} \leq \sqrt{\frac{2}{\pi}} e^{\frac{1}{2}\zeta^2} e^{-|\zeta|x} \leq \sqrt{\frac{2}{\pi}}.$$

□

REFERENCES

- [1] ATAR, R. (2012). A diffusion regime with nondegenerate slowdown. *Operations Research*, **60** 490–500. URL <http://dx.doi.org/10.1287/opre.1110.1030>.
- [2] BARBOUR, A. (1990). Stein’s method for diffusion approximations. *Probability Theory and Related Fields*, **84** 297–322. URL <http://dx.doi.org/10.1007/BF01197887>.
- [3] BARBOUR, A. D. (1988). Stein’s method and Poisson process convergence. *Journal of Applied Probability*, **25** pp. 175–184. URL <http://www.jstor.org/stable/3214155>.
- [4] BRAVERMAN, A. and DAI, J. G. (2015). Stein’s method for steady-state diffusion approximations. Preprint, URL <http://people.orie.cornell.edu/jdai/publications/preprints.html>.
- [5] CHATTERJEE, S. (2014). A short survey of Stein’s method. To appear in Proceedings of ICM 2014, URL <http://arxiv.org/abs/1404.1392>.
- [6] CHEN, L. H. Y., GOLDSTEIN, L. and SHAO, Q.-M. (2011). *Normal approximation by Stein’s method*. Probability and its Applications (New York), Springer, Heidelberg. URL <http://dx.doi.org/10.1007/978-3-642-15007-4>.
- [7] GIBBS, A. L. and SU, F. E. (2002). On choosing and bounding probability metrics. *International Statistical Review / Revue Internationale de Statistique*, **70** pp. 419–435. URL <http://www.jstor.org/stable/1403865>.
- [8] GLYNN, P. W. and ZEEVI, A. (2008). Bounding stationary expectations of Markov processes. In *Markov processes and related topics: a Festschrift for Thomas G. Kurtz*, vol. 4 of *Inst. Math. Stat. Collect.* Inst. Math. Statist., Beachwood, OH, 195–214. URL <http://dx.doi.org/10.1214/074921708000000381>.
- [9] GURVICH, I. (2014). Diffusion models and steady-state approximations for exponentially ergodic Markovian queues. *The Annals of Applied Probability*, **24** 2527–2559. URL <http://dx.doi.org/10.1214/13-AAP984>.
- [10] GURVICH, I., HUANG, J. and MANDELBAUM, A. (2014). Excursion-based universal approximations for the Erlang-A queue in steady-state. *Mathematics of Operations Research*, **39** 325–373. URL <http://dx.doi.org/10.1287/moor.2013.0606>.
- [11] HALFIN, S. and WHITT, W. (1981). Heavy-traffic limits for queues with many exponential servers. *Oper. Res.*, **29** 567–588.
- [12] HENDERSON, S. G. (1997). *Variance reduction via an approximating Markov process*. Ph.D. thesis, Department of Operations Research, Stanford University. <http://people.orie.cornell.edu/shane/pubs/thesis.pdf>.
- [13] MEYN, S. P. and TWEEDIE, R. L. (1993). Stability of Markovian processes III: Foster-Lyapunov criteria for continuous time processes. *Adv. Appl. Probab.*, **25** 518–548.

- [14] ROSS, N. (2011). Fundamentals of Stein's method. *Probab. Surv.*, **8** 210–293. URL <http://dx.doi.org/10.1214/11-PS182>.
- [15] STEIN, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*. University of California Press, Berkeley, Calif., 583–602. URL <http://projecteuclid.org/euclid.bsmmsp/1200514239>.
- [16] STEIN, C. (1986). Approximate computation of expectations. *Lecture Notes-Monograph Series*, **7**. URL <http://www.jstor.org/stable/4355512>.
- [17] WARD, A. and GLYNN, P. (2003). A diffusion approximation for a markovian queue with reneging. *Queueing Systems*, **43** 103–128. URL <http://dx.doi.org/10.1023/A:3A1021804515162>.
- [18] WHITT, W. (2003). How multiserver queues scale with growing congestion-dependent demand. *Operations Research*, **51** 531–542.

E-MAIL: ab2329@cornell.edu

E-MAIL: jd694@cornell.edu

E-MAIL: jf646@cornell.edu